Last updated: Dec 13, 2024 Abstract Algebra Notes Week 15 Wed, Dec 11 2024

Sec 14.1 Groups acting on sets & Thm 9.10 Cayley's Theorem

Intuitively, a group G "acts" on a set S of configurations by "naturally permuting" the configurations in S.

$$G = D_4 = \text{Symmetry group of a square}$$

$$= \left\{ c, R, R^2, R^3, f, fR, fR^2, fR^3 \right\} = \left\langle R, f \right\rangle$$

R = counterclockwise rotation by 90°,

f = horizontal flip

Configurations of the square mattress (corners labeled by 1,2,3,4):


Let $X = \{1, 2, 3, 4\}$, just a set.

We can think of D₄ as the following permutations in S₄:


e R R^2 R^3 f fR fR^2 fR^3 e, (1234), (13)(24), (1432), (12)(34), (13), (14)(23), (24)

$$\begin{bmatrix} 1 & 2 & \\ 4 & 3 & \\ \end{bmatrix} \begin{pmatrix} A & \\ \\ - & \\ \end{bmatrix} \begin{pmatrix} \xi & \\ \\ \\ \end{bmatrix} \begin{pmatrix} A & \\ \\ \\ \\ \\ \end{pmatrix} \begin{pmatrix} A & \\ \\ \\ \end{pmatrix} \begin{pmatrix} A & \\ \\ \\ \\ \end{pmatrix} \begin{pmatrix} A & \\ \\ \\ \\ \end{pmatrix} \begin{pmatrix} A & \\ \\ \end{pmatrix} \begin{pmatrix} A & \\ \\ \\ \end{pmatrix} \begin{pmatrix} A & \\ \\ \\ \end{pmatrix} \begin{pmatrix} A & \\ \\ \end{pmatrix} \end{pmatrix} \begin{pmatrix} A & \\ \\ \end{pmatrix} \end{pmatrix} \begin{pmatrix} A & \\ \\ \end{pmatrix} \begin{pmatrix} A & \\ \\ \end{pmatrix} \begin{pmatrix} A & \\ \\ \end{pmatrix} \end{pmatrix} \begin{pmatrix} A & \\ \\ \end{pmatrix} \begin{pmatrix} A & \\ \\ \end{pmatrix} \begin{pmatrix} A & \\ \\ \end{pmatrix} \end{pmatrix} \begin{pmatrix} A & \\ \\ \end{pmatrix} \begin{pmatrix} A & \\ \\ \end{pmatrix} \begin{pmatrix} A & \\ \\ \end{pmatrix} \end{pmatrix} \begin{pmatrix} A & \\ \\ \end{pmatrix} \begin{pmatrix} A & \\ \\ \end{pmatrix} \end{pmatrix} \begin{pmatrix} A & \\ \\ \end{pmatrix} \begin{pmatrix} A & \\ \\ \end{pmatrix} \end{pmatrix} \begin{pmatrix} A & \\ \\ \end{pmatrix} \begin{pmatrix} A & \\ \\ \end{pmatrix} \end{pmatrix} \begin{pmatrix} A & \\ \\ \end{pmatrix} \begin{pmatrix} A & \\ \\ \end{pmatrix} \end{pmatrix} \begin{pmatrix} A & \\ \\ \end{pmatrix} \begin{pmatrix} A & \\ \\ \end{pmatrix} \end{pmatrix} \begin{pmatrix} A & \\ \\ \end{pmatrix} \begin{pmatrix} A & \\ \\ \end{pmatrix} \end{pmatrix} \begin{pmatrix} A & \\ \\ \end{pmatrix} \begin{pmatrix} A & \\ \\ \end{pmatrix} \end{pmatrix} \begin{pmatrix} A & \\ \\ \end{pmatrix} \begin{pmatrix} A & \\ \\ \end{pmatrix} \end{pmatrix} \begin{pmatrix} A & \\ \\ \end{pmatrix} \begin{pmatrix} A & \\ \\ \end{pmatrix} \end{pmatrix} \begin{pmatrix}$$

Action diagram:



Note: This diagram is connected



Note: This diagram is not connected

Remark (Extra): When the action diagram is connected, the group action is called <u>transitive</u>

(from one elt a in the set X, we can get to all other elts in X using G)

where

1.
$$ex = x$$
 for all $x \in X$

2.
$$(ba)x = b(ax)$$
 for all $x \in X$ and $a, b \in G$.

A set X equipped w/ such a map is called a left G-set.

Note: X doesn't need to be related to G in any way. But group actions are more interesting when the G-sets X is related to the group G in some way.

Ex 3 (Ex of group actions from linear algebra)

 $(t \times 14.1)$

 $G = GL_2(R) = \{2x2 \text{ invertible matrices } w \text{ real entries}\}$ $\overline{X} = R^2 = \{\text{ vectors } \begin{bmatrix} x \\ y \end{bmatrix} : x, y \in R \}$

Then G acts on X by left multiplication $GL_{2}(\mathbb{R}) \times \mathbb{R}^{2} \longrightarrow \mathbb{R}^{2}$

$$\left(\begin{bmatrix} a & b \\ c & d \end{bmatrix}, \quad \begin{bmatrix} x \\ y \end{bmatrix} \right) \mapsto \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

Verify that this map satisfies the two conditions:

1.
$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x \\ y \end{bmatrix}$$
 for all $\begin{bmatrix} x \\ y \end{bmatrix} \in \mathbb{R}^2$

2. For all A, B & GL2 (R), we have

$$(BA)\begin{bmatrix} x \\ y \end{bmatrix} = B(A\begin{bmatrix} x \\ y \end{bmatrix})$$

since matrix multiplication is associative.

Ex I and 2 are group actions for D4

Group guiz will be to rewrite Ex 3 using alt def

Def Perm (X) = the group of permutations of X bijections $X \to X$ Note: If |X| = n then Perm (X) \cong Sn

An alternative definition of group action (equivalent to the defin the book)

Intuition

- · Given a group G, we have a "switch board" with a button 9 for every g & G.
- Given $a \in G$, pressing the button a rearranges the objects in our set X. This gives a permutation on X; call this $p(a) \in Perm(X)$
- Given b \in G, pressing the button \bigcirc also rearranges the objects in our set \boxtimes . Call this permutation ϕ (b).
- · The element ba & G also has a button ba
- For G to act on X, we require that pressing the ba button gives the same rearrangement of X as first pressing the Dutton, followed by the button \Box , that is, $\frac{d}{dt} \left(\frac{dt}{dt} \right) = \frac{d}{dt} \left(\frac{dt}{dt}$

$$\phi(ba) = \phi(b) \phi(a)$$

for all a, b∈ G


Alternative def of group action

A <u>left action</u> of G on X is a group homomorphism $\phi: G \longrightarrow \text{Perm}(X)$


Note: A right action requires \$\phi(ba) = \phi(a) \phi(b) instead of (*)

Back to Ex 1

"Switchboard" for D4;




Pressing the R button permutes X = {1,2,3,4} as follows:



$$\phi(R) = (1234)$$

Pressing the f button permutes X = {1,2,3,4} as follows:



$$\phi$$
 (D4) = { e, (1234), (13)(24), (1432), (12)(34), (13), (14)(23), (24) }
 $\leq S_4$
Subgroup

Group quit: In Example 2, think of the xxoo squares as 1,2,...,6. Write the left action of Dq on {1,2,...6} Write the elts of im p. Is it isomorphic to a familiar group?

Next, we see that every group acts on itself by left multiplication.

Def Let X = G. Define the homomorphism $(E \times 14.3)$

$$\phi: G \longrightarrow \overline{Perm(G)}$$
 by $g \longmapsto T_{\mathfrak{g}}$

where

$$T_g: G \to G$$

 $\times \mapsto g \times \text{ for all } \times \in G$ (that is, T_g is multiplication)
by g on the left

In other words, pressing the g button on our "group switchboard" multiplies every elt on the left by g.

The image $\phi(G) = \{T_g : g \in G\}$ is called the left regular representation of G

Recall:
The units in
Z12 are
1,5,7,11,&
They form
a group wy
operation.

Exercise: Prove that & is a group homomorphism

Prove that \$\phi\$ is injective

Ex Calculate the left originar rep of U(12)= {1,5,7,11}

$$T_1 = multp \ by \ 1 = \begin{bmatrix} 1 & 5 & 7 & 11 \\ 1 & 5 & 7 & 11 \end{bmatrix}$$

$$T_5 = multp \ by \ 5 = \begin{bmatrix} 1 & 5 & 7 & 11 \\ 5 & 1 & 11 & 7 \end{bmatrix}$$

$$= (ab) (cd)$$

$$T_{7} = \text{multp by } 7 = \begin{bmatrix} 1 & 5 & 7 & 11 \\ 7 & 11 & 1 & 5 \end{bmatrix}$$

$$T_{11} = \begin{bmatrix} 1 & 5 & 7 & 11 \\ 11 & 7 & 5 & 1 \end{bmatrix}$$

$$q_{11} = \begin{bmatrix} 1 & 7 & 11 \\ 11 & 7 & 5 & 1 \end{bmatrix}$$

$$q_{11} = \begin{bmatrix} 1 & 5 & 7 & 11 \\ 11 & 7 & 5 & 1 \end{bmatrix}$$

$$q_{11} = \begin{bmatrix} 1 & 5 & 7 & 11 \\ 11 & 7 & 5 & 1 \end{bmatrix}$$

$$q_{11} = \begin{bmatrix} 1 & 5 & 7 & 11 \\ 11 & 7 & 5 & 1 \end{bmatrix}$$

Note (extra) Every group also acts on itself by right multiplication. The corresponding action diagram is equal to the Cayley diagram.

Cayley's Thm Every group is isomorphic to (See Thm 9.12) a group of permutations.

Pf Let ϕ be the left regular representation $\phi: G \longrightarrow \operatorname{Perm}(G)$ $g \longmapsto \operatorname{Tg}$

given above.

Then in ϕ is group of permutations. Since ϕ is injective, we have $G \cong \text{im } \phi$. \square

Two contrasting reasons why Cayley's Thom is important

- (1) Cayley's Thm allows us to represent an abstract group in a concrete way
- 2) Present-day set of axioms for a group is
 the correct abstraction of a group of permutations.

Note: Concepts similar to "group acting on set" appear in other algebraic structures.

- · For example, in ring theory, we have "R-modules" instead of "G-sets".
- · If you are interested in learning about module theory next semester / year, come talk to me! ______ end ___