

Math 3230 Abstract Algebra I

Sec 3.2: Cosets

Slides created by M. Macauley, Clemson (Modified by E. Gunawan, UConn)


`http://egunawan.github.io/algebra`

Abstract Algebra I

Idea of cosets

Copies of the fragment of the Cayley diagram that corresponds to a subgroup appear throughout the rest of the diagram.

Example: Below you see three copies of the fragment corresponding to the subgroup $\langle f \rangle = \{e, f\}$ in D_3 .



However, only one of these copies is actually a group! Since the other two copies do *not* contain the identity, they cannot be groups.

Key concept

The elements that form these repeated copies of the fragment of a subgroup H in the Cayley diagram are called **cosets of H** .

Above show the three cosets of the subgroup $\{e, f\}$.

An example: D_4



Let $H = \langle f, r^2 \rangle = \{e, f, r^2, r^2f\}$, a subgroup of D_4 . Find all of the cosets of H .

If we use r^2 as a generator in the Cayley diagram of D_4 , then it will be easier to “see” the cosets.

Note that $D_4 = \langle r, f \rangle = \langle r, f, r^2 \rangle$. The cosets of $H = \langle f, r^2 \rangle$ are:

$$H = \langle f, r^2 \rangle = \underbrace{\{e, f, r^2, r^2f\}}_{\text{original}}$$

$$rH = r\langle f, r^2 \rangle = \underbrace{\{r, r^3, rf, r^3f\}}_{\text{copy}}$$



Definition of cosets

Definition

If H is a subgroup of G , then a (left) **coset** of H is a set

$$xH = \{xh : h \in H\},$$

where $x \in G$ is some fixed element. The distinguished element (in this case, x) that we choose to use to name the coset is called the **representative**.

Remark


In a Cayley diagram, the (left) coset xH can be found as follows: **start from node x and follow all paths in H .**

For example, let $H = \langle f \rangle$ in D_3 . The coset $\{r, rf\}$ of H is the set

$$rH = r\langle f \rangle = r\{e, f\} = \{r, rf\}.$$

Alternatively, we could have written $(rf)H$ to denote the same coset, because

$$rfH = rf\{e, f\} = \{rf, rf^2\} = \{rf, r\}.$$



More on cosets

Proposition 1

For any subgroup $H \leq G$, the union of the (left) cosets of H is the whole group G .

Proof

We only need to show that every element $x \in G$ lives in some coset of H . But, since $e \in H$ (because H is a group) and $x = xe$, we can conclude that x lives in the coset $xH = \{xh \mid h \in H\}$. \square

Proposition 2 (HW)

If $y \in xH$, then $xH = yH$. \square

More on cosets

Proposition 3 (HW)

All (left) cosets of a subgroup H of G have the same size as H . □

Hint: Define a bijection between $eH = H$ and another coset xH . Copy the bijection between the even permutations and odd permutations from notes 2.4, but replace (12) with x .

More on cosets

Proposition 4

For any subgroup $H \leq G$, the (left) cosets of H **partition** the group G .

Proof

To show that the set of (left) cosets of H form a partition of G , we need to show that (1) the union of all (left) cosets of H is equal to G , and (2) if H is a proper subgroup, then the intersection of each pair of two distinct (left) cosets of H is empty.

Part (1) has been shown earlier in Proposition 1: every element x is the coset xH . To show part (2), suppose that $x \in G$ lies in a coset yH . Then by Proposition 2, $xH = yH$. So every element of G lives in exactly one coset. □

Subgroups also have **right cosets**: $Ha = \{ha : h \in H\}$.

For example, the three right cosets of $H = \langle f \rangle$ in D_3 are H ,


$$\begin{aligned} Hr &= \langle f \rangle r = \{e, f\}r = \{r, fr = r^2f\}, \text{ and} \\ \langle f \rangle r^2 &= \{e, f\}r^2 = \{r^2, fr^2\} = \{r^2, rf\}. \end{aligned}$$

In this example, the left cosets for $\langle f \rangle$ are **different** from the right cosets.

Left vs. right cosets

The left diagram below shows the **left coset** $r\langle f \rangle$ in D_3 : the nodes that f arrows can reach **after** the path to r has been followed.

The right diagram shows the **right coset** $\langle f \rangle r$ in D_3 : the nodes that r arrows can reach **from** the elements in $\langle f \rangle$.



Left cosets look like copies of the subgroup, while the elements of right cosets are usually scattered (only because we adopted the convention that arrows in a Cayley diagram represent **right multiplication**).

Key point




Left and right cosets are generally different.

Left vs. right cosets

For *any* subgroup $H \leq G$, we can think of G as the union of non-overlapping and equal size copies of H , namely the left cosets of H .

Though the right cosets also partition G , the corresponding partitions could be different!

Here are a few visualizations of this idea:



Definition

If $H < G$, then the **index** of H in G , written $[G : H]$, is the number of distinct left (or equivalently, right) cosets of H in G .

Left vs. right cosets

The left and right cosets of the subgroup $H = \langle f \rangle \leq D_3$ are *different*:

| | | | | | | | | |
|--------|----------------------------------------------------------------------------------------|--------|-------|--|----------------------------------------------------------------------------------------|--------|-------|--|
| r^2H | <table border="1"><tr><td>r^2f</td><td>r^2</td></tr></table> | r^2f | r^2 | | <table border="1"><tr><td>r^2f</td><td>r^2</td></tr></table> | r^2f | r^2 | |
| r^2f | r^2 | | | | | | | |
| r^2f | r^2 | | | | | | | |
| rH | <table border="1"><tr><td>r</td><td>rf</td></tr></table> | r | rf | | <table border="1"><tr><td>r</td><td>rf</td></tr></table> | r | rf | |
| r | rf | | | | | | | |
| r | rf | | | | | | | |
| H | <table border="1"><tr><td>e</td><td>f</td></tr></table> | e | f | | <table border="1"><tr><td>e</td><td>f</td></tr></table> | e | f | |
| e | f | | | | | | | |
| e | f | | | | | | | |

| | | | | | | |
|--------|-------|------|----------------------------------------------------------------------------------------|--------|-------|--------|
| | | Hr | <table border="1"><tr><td>r^2f</td><td>r^2</td></tr></table> | r^2f | r^2 | |
| r^2f | r^2 | | | | | |
| | | | <table border="1"><tr><td>r</td><td>rf</td></tr></table> | r | rf | Hr^2 |
| r | rf | | | | | |
| | | H | <table border="1"><tr><td>e</td><td>f</td></tr></table> | e | f | |
| e | f | | | | | |

The left and right cosets of the subgroup $N = \langle r \rangle \leq D_3$ are *the same*:

| | | | | | | | | | | |
|------|--------------------------------------------------------------------------------------------------------------|--------|------|--------|--|------|--------------------------------------------------------------------------------------------------------------|-----|------|--------|
| fN | <table border="1"><tr><td>f</td><td>rf</td><td>r^2f</td></tr></table> | f | rf | r^2f | | Nf | <table border="1"><tr><td>f</td><td>rf</td><td>r^2f</td></tr></table> | f | rf | r^2f |
| f | rf | r^2f | | | | | | | | |
| f | rf | r^2f | | | | | | | | |
| N | <table border="1"><tr><td>e</td><td>r</td><td>r^2</td></tr></table> | e | r | r^2 | | N | <table border="1"><tr><td>e</td><td>r</td><td>r^2</td></tr></table> | e | r | r^2 |
| e | r | r^2 | | | | | | | | |
| e | r | r^2 | | | | | | | | |

Proposition 5 (HW)

If $H \leq G$ has index $[G : H] = 2$, then the left and right cosets of H are the same.

Cosets of abelian groups

Recall: in some abelian groups, we use the symbol $+$ for the binary operation.

In this case, we write the left cosets as $a + H$ (instead of aH).

For example, let $G := (\mathbb{Z}, +)$, and consider the subgroup $H := 4\mathbb{Z} = \{4k \mid k \in \mathbb{Z}\}$ of G consisting of multiples of 4. Then the left cosets of H are

$$\begin{aligned}H &= \{\dots, -12, -8, -4, 0, 4, 8, 12, \dots\} \\1 + H &= \{\dots, -11, -7, -3, 1, 5, 9, 13, \dots\} \\2 + H &= \{\dots, -10, -6, -2, 2, 6, 10, 14, \dots\} \\3 + H &= \{\dots, -9, -5, -1, 3, 7, 11, 15, \dots\}.\end{aligned}$$

Notice that these are the same as the right cosets of H :

$$H, \quad H + 1, \quad H + 2, \quad H + 3.$$

Exercise: Why are the left and right cosets of an abelian group *always* the same?

Note that it would be confusing to write $3H$ for the coset $3 + H$. In fact, $3H$ would usually be interpreted to mean the subgroup $3(4\mathbb{Z}) = 12\mathbb{Z}$.

A theorem of Joseph Lagrange

Lagrange's Theorem

Assume G is finite. If $H < G$, then $|H|$ divides $|G|$.

Proof

Suppose there are n left cosets of the subgroup H . Since they are all the same size (by Proposition 3) and they partition G (by Proposition 4), we must have

$$|G| = \underbrace{|H| + \cdots + |H|}_{n \text{ copies}} = n|H|.$$

Therefore, $|H|$ divides $|G|$. □

Corollary of Lagrange's Theorem

If G is a finite group and $H \leq G$, then

$$[G : H] = \frac{|G|}{|H|}.$$

A theorem of Joseph Lagrange


Corollary of Lagrange's Theorem

If G is a finite group and $H \leq G$, then $[G : H] = \frac{|G|}{|H|}$

This significantly narrows down the possibilities for subgroups.

Warning: The converse of Lagrange's Theorem is not generally true. That is, just because $|G|$ has a divisor d does *not* mean that there is a subgroup of order d .

The subgroup lattice of D_4 :



From HW: Find all subgroups of $S_3 = \{e, (12), (23), (13), (123), (132)\}$ and arrange them in a subgroup lattice.