

# Math 3230 Abstract Algebra I

## Sec 4.5: Isomorphism theorems

Slides created by M. Macauley, Clemson (Modified by E. Gunawan, UConn)

`http://egunawan.github.io/algebra`

Abstract Algebra I

## The Isomorphism Theorems

The Fundamental Homomorphism Theorem (FHT) is the first of four basic theorems about homomorphism and their structure.

These are commonly called “**The Isomorphism Theorems**”:

- First Isomorphism Theorem: “Fundamental Homomorphism Theorem”
- Second Isomorphism Theorem: “Diamond Isomorphism Theorem”
- Third Isomorphism Theorem: “Freshman Theorem”
- Fourth Isomorphism Theorem: “Correspondence Theorem”

All of these theorems have analogues in other algebraic structures: rings, vector spaces, modules, and Lie algebras, to name a few.

In this lecture, we will summarize the last three isomorphism theorems and provide visual pictures for each.

We will prove one, outline the proof of another (homework!), and encourage you to try the (very straightforward) proofs of the multiple parts of the last one.

Finally, we will introduce the concepts of a **commutator** and **commutator subgroup**, whose quotient yields the **abelianization** of a group.


# The Second Isomorphism Theorem

## Diamond isomorphism theorem

Let  $H < G$ , and  $N \triangleleft G$ . Then

- (i) The **product**  $HN = \{hn \mid h \in H, n \in N\}$  is a subgroup of  $G$ .
- (ii) The **intersection**  $H \cap N$  is a *normal* subgroup of  $H$ .
- (iii) The following quotient groups are isomorphic:

$$HN/N \cong H/(H \cap N)$$



## Proof (sketch)

Define the following map

$$\phi: H \longrightarrow HN/N, \quad \phi: h \longmapsto hN.$$

If we can show:

1.  $\phi$  is a homomorphism,
2.  $\phi$  is surjective (onto),
3.  $\text{Ker } \phi = H \cap N$ ,

then the result will follow *immediately* from the FHT. Watch M. Macauley's video of Slides 4.5 and Judson's textbook Theorem 11.12 for more details. The rest is left as exercise.

# The Third Isomorphism Theorem


## Freshman theorem

Consider a chain  $N \leq H \leq G$  of normal subgroups of  $G$ .

That is, both  $N$  and  $H$  are normal subgroups of  $G$ , and  $N < H$ . Then

1. The quotient  $H/N$  is a normal subgroup of  $G/N$ ;
2. The following quotients are isomorphic:

$$(G/N)/(H/N) \cong G/H.$$



(Concept and graphics by Zach Teitler of Boise State)

# The Third Isomorphism Theorem

## Freshman theorem

Consider a chain  $N \leq H \leq G$  of normal subgroups of  $G$ . Then (i)  $H/N \trianglelefteq G/N$  and (ii)  $(G/N)/(H/N) \cong G/H$ .

## Proof

(i) The fact that  $H/N \trianglelefteq G/N$  is shown in M. Macauley's video. To show (ii), define the map

$$\varphi: G/N \longrightarrow G/H, \quad \varphi: gN \longmapsto gH.$$

- Show  $\varphi$  is well-defined: Suppose  $g_1N = g_2N$ . Then  $g_1 = g_2n$  for some  $n \in N$ . But  $n \in H$  because  $N \leq H$ . Thus,  $g_1H = g_2H$ , i.e.,  $\varphi(g_1N) = \varphi(g_2N)$ . ✓
- Show that  $\varphi$  is onto and a homomorphism. ✓
- Apply the FHT:

$$\begin{aligned} \text{Ker } \varphi &= \{gN \in G/N \mid \varphi(gN) = H\} \\ &= \{gN \in G/N \mid gH = H\} \\ &= \{gN \in G/N \mid g \in H\} = H/N \end{aligned}$$

By the FHT,  $(G/N)/\text{Ker } \varphi = (G/N)/(H/N) \cong \text{Im } \varphi = G/H$ . □




# The Fourth Isomorphism Theorem

## Correspondence theorem (idea)

Let  $N \triangleleft G$ . There is a 1-1 correspondence between **subgroups of  $G/N$**  and **subgroups of  $G$  that contain  $N$** . In particular, every subgroup of  $G/N$  can be written as  $\bar{A} := A/N$  for some  $A$  satisfying  $N < A < G$ .

This means that the corresponding subgroup lattices are identical in structure.

## Example



Here  $G = Q_8$  and  $N = \{1, -1\}$ .




The quotient  $Q_8 / \langle -1 \rangle$  is isomorphic to  $V_4$ . The subgroup lattices can be visualized by “collapsing”  $\langle -1 \rangle$  to the identity.

## Correspondence theorem (formally)

Let  $N \triangleleft G$ . Then there is a bijection from the **subgroups of  $G/N$**  and **subgroups of  $G$  that contain  $N$** . In particular, every subgroup of  $G/N$  has the form  $\bar{A} := A/N$  for some  $A$  satisfying  $N < A < G$ . Moreover, if  $A, B < G$ , then

1.  $A \leq B$  if and only if  $\bar{A} \leq \bar{B}$ ,
2. If  $A \leq B$ , then  $[B : A] = [\bar{B} : \bar{A}]$ ,
3.  $\overline{\langle A, B \rangle} = \langle \bar{A}, \bar{B} \rangle$ ,
4.  $\overline{A \cap B} = \bar{A} \cap \bar{B}$ ,
5.  $A \triangleleft G$  if and only if  $\bar{A} \triangleleft \bar{G}$ .

## Example with $G = D_4$ and $N = \langle r^2 \rangle$



## Application: commutator subgroups and abelianizations



Idea: We've seen how to divide  $\mathbb{Z}$  by  $12\mathbb{Z}$ , thereby "forcing" all multiples of 12 to be zero. This is one way to construct the integers modulo 12:  $\mathbb{Z}_{12} \cong \mathbb{Z}/12\mathbb{Z}$ .

Now, let  $G$  be nonabelian. We want to divide  $G$  by its "non-abelian parts," making them zero and leaving only "abelian parts" in the resulting quotient.

### Definition

A **commutator** of a group  $G$  is an element of the form  $aba^{-1}b^{-1}$ , where  $a, b \in G$ .

If  $G$  is nonabelian, there are non-identity commutators:  $aba^{-1}b^{-1} \neq e$  in  $G$ .



In this case, the set  $C := \{aba^{-1}b^{-1} \mid a, b \in G\}$  contains *more* than the identity.

### Definition

Define the **commutator subgroup**  $G'$  of  $G$  to be  $G' := \langle aba^{-1}b^{-1} \mid a, b \in G \rangle$ .

This is a normal subgroup of  $G$  (exercise). If we quotient out by it, we get an abelian group! (Because we have killed every instance of the " $ab \neq ba$  pattern")



## Commutator subgroups and abelianizations

### Definition


The **abelianization** of  $G$  is the quotient group  $G/G'$ . This is the group that one gets by “killing off” all nonabelian parts of  $G$ .

In some sense, the commutator subgroup  $G'$  is the **smallest normal subgroup**  $N$  of  $G$  such that  $G/N$  is abelian. [Note that  $G$  would be the “largest” such subgroup.]

Equivalently, the quotient  $G/G'$  is the **largest abelian quotient** of  $G$ . [Note that  $G/G \cong \{e\}$  would be the “smallest” such quotient.]

### Universal property of commutator subgroups

Suppose  $f: G \rightarrow A$  is a homomorphism to an abelian group  $A$ . Then there is a unique homomorphism  $h: G/G' \rightarrow A$  such that  $f = hq$ :





We say that  $f$  “factors through” the abelianization,  $G/G'$ .

# Commutator subgroups and abelianizations

## Examples

Check that the commutator subgroups of  $A_4$  and  $D_4$  are

$$A_4' = \langle xyx^{-1}y^{-1} \mid x, y \in A_4 \rangle \cong V_4, \quad D_4' = \langle xyx^{-1}y^{-1} \mid x, y \in D_4 \rangle = \langle r^2 \rangle.$$



By the *Correspondence Theorem*, the abelianization of  $A_4$  is  $A_4/V_4 \cong C_3$ , and the abelianization of  $D_4$  is  $D_4/\langle r^2 \rangle \cong V_4$ .

Notice that  $G/G'$  is abelian, and moreover, taking the quotient of  $G$  by *any subgroup drawn above  $G'$*  will yield an abelian group.