

Last updated on 2016/12/13 at 13:27:30. Let me know of typos.

1 Mostly topics and skills from Test 1

1.1 Test 1: slope, secant line, tangent line, limit, derivative

1. Let $F(t)$ be a function defined by $F(t) = t^2 + 3t - 1$.

(a) Evaluate the *difference quotient*

$$\frac{F(t + \Delta t) - F(t)}{\Delta t}.$$

(b) What is the slope of the secant line of the graph of $F(t)$ over $[t, t + \Delta t]$

(c) What is the slope of the tangent line of the graph of $F(t)$ at t ?

(d) Compute $F'(t)$.

(e) Compute the *net change* in F over the interval $[-1, 4]$.

(f) Compute the *average rate of change* in F over the interval $[-1, 4]$.

(g) Write an equation for the secant line which meets the graph of f at $x = -1$ and $x = 4$.

(h) Write an equation for the secant line which meets the graph of f at $x = a$ and $x = 4$, where a is a real number that is smaller than 4.

2. Let $f(x) = x^3 - 2$.

(a) Evaluate $f(x + \Delta x)$ (your answer should be a simple formula involving variables x and Δx).

(b) Evaluate and simplify the following quotient as much as possible.

$$\frac{f(x + \Delta x) - f(x)}{\Delta x}.$$

(c) Compute $\lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x}$.

(d) Use the previous answer to compute $f'(x)$.

3. Function g is defined by

$$g(x) = \frac{(x - 4)^{999} + 2x}{70}.$$

(a) $g(5) - g(4)$

(b) What is the *average rate of change* in g over the interval $[4, 5]$?

(c) Write an equation for the secant line which meets the graph of g at $x = 4$ and $x = 5$.

(d) $g(x + 4)$.

1.2 Test 1: Some Concepts from Test 1


1. True or False: If the average rate of change in a function f over $[a, b]$ is positive, then f must be increasing over $[a, b]$.
2. T/F: If f is a linear function, then f has the same net change over every possible interval.
3. T/F: If f is a linear function, then f has the same average rate of change over every possible interval.
4. Which of the following functions has/ have a constant rate of change (the same average rate of change over every possible interval)?
 - a. $\frac{1}{9}x$
 - b. $\frac{9}{x}$
 - c. $\frac{1}{9x}$
 - d. $(\frac{1}{9}x - 1)(x + 9)$
 - e. $9x - \frac{1}{9}$

1.3 Test 1: Finding zeros of polynomials

1. Find the zeros of the following functions. If no zeros exist, say so.
 - (a) $f(t) = t^{10} - 2t^9$.
 - (b) $h(x) = (x - 2)(x^2 + 1)$.
 - (c) $g(x) = (x - 5)(x^3 + 1)$. (Note: this will involve long division).
 - (d) $j(x) = (x + 5)^3(x^2 - 2)(x - \frac{1}{2})$. Note that $x^2 - 2 = (x - \sqrt{2})(x + \sqrt{2})$, so $j(x) = (x + 5)^3(x - \sqrt{2})(x + \sqrt{2})(x - \frac{1}{2})$.
 - (e) $h(x) = x^2 + x + 3$.

1.4 Test 1: More Skills from Test 1

The figure shows the graph of a quadratic function:



1. Find the *exact* coordinates of each of the three marked points A , B , and C . Note, A is the point where the graph crosses the y -axis; B and C are the points where it crosses the x -axis.
2. Write down exactly all the *zero/s* (aka *root/s*) of this function.
3. What degree is this polynomial? What is the leading term of this polynomial?
4. How many relative max or relative min does this graph has?
5. Write an equation of the secant line between the points $(0, 3)$ and $(2, 5)$.
6. Write an equation of the secant line between the points $(1, 6)$ and $(2, 5)$.

1.5 Test 1: End behaviors and zeros of polynomials

1. Give an example of a polynomial of degree 7 which falls to the left, rises to the right, and has $x=2$, $x=5$, and $x=11$ as zeros. You do not need to sketch the graph, but you need to write the formula for the polynomial.
2. Give an example of a polynomial of degree 5 which rises to the left, falls to the right, and has $x=2$ as a zero. You do not need to sketch the graph, but you need to write the formula for the polynomial.
3. Give an example of a polynomial of degree 6 which falls to the left, falls to the right, and has $x=2$ as a zero. You do not need to sketch the graph, but you need to write the formula for the polynomial.
4. Give an example of a polynomial which has $x = 2$ as a zero but does *not* change sign at $x = 2$.
5. Give an example of a polynomial which has $x = 2$ as a zero and changes sign from positive to negative at $x = 2$.
6. True or False: every polynomial of even degree (2 or higher) has at least a zero. (If you write True, give an explanation. If write False, give a counterexample. If you give a counterexample, you need to give the formula for the polynomial.)
7. True or False: the polynomial $f(x) = x^2 - 4$ has exactly two zeros,
8. True or False: the polynomial $f(x) = x^2 - 4$ has a vertical asymptote at $x = -2$.
9. True or False: every polynomial of odd degree has at least one zero.

2 From Test 2

2.1 Test 2: Evaluating limits

1. Let $F(t)$ be a function defined by

$$F(t) = \frac{t^2 + 3t - 1}{5}.$$

- (a) Evaluate $\lim_{t \rightarrow 0} F(t)$
 - (b) Evaluate $\lim_{t \rightarrow 1} F(t)$
 - (c) Identify any points at which F is discontinuous.
2. Evaluate $\lim_{t \rightarrow 0} 7 + x - \sqrt{7} + \frac{x+2}{x+1}$.
 3. Evaluate $\lim_{t \rightarrow 0^+} \frac{\sqrt{7+x} - \sqrt{7-x}}{2x}$.
 4. Evaluate $\lim_{t \rightarrow 1^-} \frac{\sqrt{2x+1} - \sqrt{3}}{x-1}$.

2.2 Test 2: Identifying Vertical asymptotes analytically

Identify the vertical asymptotes (if any) of following functions. If the function has no vertical asymptotes, say so.

1.

$$f(t) = \frac{5}{t^{10} - 2t^9}.$$


2.

$$g(x) = \frac{x+1}{(x-5)(x^3+1)}.$$

3. $h(x) = x^2 - 25x^2 + x + 3$.

2.3 Test 2: Reading and sketching graphs

The figure shows the graph of a function $f(x)$.



Read the value graph; if it does not exist or is not defined, just say so.

1. $\lim_{x \rightarrow 0^-} f(x)$

2. $\lim_{x \rightarrow 0^+} f(x)$

3. $\lim_{x \rightarrow 0} f(x)$

4. $\lim_{x \rightarrow 3^-} f(x)$

5. $\lim_{x \rightarrow 3^+} f(x)$

6. $\lim_{x \rightarrow 3} f(x)$

7. $\lim_{x \rightarrow 4^-} f(x)$

8. $\lim_{x \rightarrow 4^+} f(x)$

9. $\lim_{x \rightarrow 4} \frac{f(x)}{11}$

10. $\lim_{x \rightarrow 5^-} f(x)$

11. $\lim_{x \rightarrow 5^+} f(x)$

12. $\lim_{x \rightarrow 5} \frac{f(x)}{x}$

13. $\lim_{x \rightarrow 7^-} f(x)$

14. $\lim_{x \rightarrow 7^+} f(x)$

15. $\lim_{x \rightarrow 7} \frac{x+1}{f(x)}$

16. $f(0)$

17. $f(3)$

18. $f(4)$

19. $f(5)$

20. $f(6)$

21. $f(7)$

22. For each part, answer True / False

(a) f is continuous on $[0,2]$.(b) f is continuous on $(0,2)$.(c) f is continuous on $[1,3]$.(d) f is continuous on $(1,3)$.(e) f is continuous on $[3,5]$.(f) f is continuous on $(3,5)$.(g) f is continuous on $[3,6]$.(h) f is continuous on $(3,6)$.(i) f is continuous on $[4,6]$.(j) f is continuous on $(4,6)$.

23. For each part, answer True / False

(a) $\lim_{x \rightarrow 0^-} f(x) = f(0)$.(b) f is continuous at 0.(c) f has a removable discontinuity at 0.(d) f has a non-removable discontinuity at 0.(e) $\lim_{x \rightarrow 2^+} f(x) = f(2)$.(f) f is continuous at 2.(g) f has a removable discontinuity at 2.(h) f has a non-removable discontinuity at 2.

- (i) $\lim_{x \rightarrow 3^-} f(x) = f(3)$.
 - (j) f is continuous at 3.
 - (k) f has a removable discontinuity at 3.
 - (l) f has a non-removable discontinuity at 3.
 - (m) $\lim_{x \rightarrow 5} f(x) = f(5)$.
 - (n) f is continuous at 5.
 - (o) f has a removable discontinuity at 5.
 - (p) f has a non-removable discontinuity at 5.
 - (q) f is continuous on $(1, 3)$.
 - (r) f is continuous on $[3, 5]$.
 - (s) f is continuous on $(3, 5)$.
 - (t) f is continuous on $[3, 6]$.
 - (u) f is continuous on $(3, 6)$.
 - (v) f is continuous on $[4, 6]$.
 - (w) f is continuous on $(4, 6)$.
24. For each part, set up suitable axes and sketch the graph of a function satisfying all the given conditions. Make your drawing as clear as possible. If I cannot tell whether it's correct, I'll assume it's not.
- (a) $f(1)=2$; f has a removable discontinuity at 1, but f is continuous everywhere else.
 - (b) $f(1)$ is not defined; f has a removable discontinuity at 1, but f is continuous everywhere else.
 - (c) $f(1)=2$; f has a non-removable discontinuity at 1, but f is continuous everywhere else.
 - (d) $f(1)$ is not defined; f has a non-removable discontinuity at 1, but f is continuous everywhere else.
 - (e) f is continuous everywhere except at $x = 2$ and $x = 5$, and $\lim_{x \rightarrow 2} f(x) = +\infty$ and $\lim_{x \rightarrow 5} f(x) = -\infty$.
 - (f) f is continuous on $(2, 5)$, and $\lim_{x \rightarrow 2^+} f(x) = +\infty$ and $\lim_{x \rightarrow 5^-} f(x) = +\infty$.
 - (g) f is continuous on $(2, 5)$, and $\lim_{x \rightarrow 2^+} f(x) = +\infty$ and $\lim_{x \rightarrow 5^-} f(x) = -\infty$.
 - (h) f is continuous on $(2, 5)$, and $\lim_{x \rightarrow 2^+} f(x) = -\infty$ and $\lim_{x \rightarrow 5^-} f(x) = -\infty$.

2.4 Test 2: Intermediate Value Theorem

1. Fill each blank with either an English word or a mathematical symbol so that the following is a complete and accurate statement of the Intermediate Value Theorem.

If a _____ $f(x)$ is _____ on the _____ interval $[a, b]$, and a number M is between _____ and _____, then there is at least one point c in _____ where _____ = _____.

2. For each part, answer either True or False. For all parts, assume that f is continuous on $[0, 2]$, that $f(0) = -10$ and $f(2) = 10$.
- (a) There must be at least one value c in $(0, 2)$ where $f(c) = 10$.
 - (b) There must be at least one value c in $[0, 2]$ where $f(c) = 10$.
 - (c) There must be at least one value c in $[0, 2]$ where $f(c) = -1$.
 - (d) There must be at least one value c in $[0, 2]$ where $f(c) = -20$.
 - (e) $f(1)$ must be between -10 and 10 .

2.5 Test 2: Evaluating limits (part 2)

Evaluate the following limits. Whenever appropriate, answer with $+\infty$ or $-\infty$.

1. $\lim_{x \rightarrow 2^+} \frac{x}{x-2}$.
2. $\lim_{x \rightarrow 2} \frac{1-x}{(x-2)^2}$.
3. $\lim_{x \rightarrow 5^-} \frac{5-x}{(x-2)^2}$.
4. $\lim_{x \rightarrow 5^+} \frac{5+x}{(x-2)^2}$.
5. $\lim_{x \rightarrow -1^+} \frac{x^3+1}{x+1}$.

3 Test 3

3.1 Test 3: the derivative function as a limit

Given the following functions for $f(x)$, use the *definition* (limit in sec 4.1) of derivative to evaluate $f'(x)$.

1. $f(x) = \sqrt{x+5}$
2. $f(x) = \sqrt{x+3}$
3. $f(x) = \sqrt{x+1}$
4. $f(x) = 2x^3 + 1$

3.2 Test 3: Tangent lines

Suppose that $f(x) = \frac{-7}{(2x-3)^2}$.

1. Compute $f'(x)$ using derivative rules.
2. Find the slope of the tangent line to the graph of f at $x = 5$
3. Write an equation for the tangent line at $x = 5$.

3.3 Test 3: Applying the power rule, constant multiple, product rule, and quotient rule. Plus graphical applications of derivatives.

Compute the derivative of each of the following functions. Use the derivative rules. Do not set up a limit for these.

1. How do you explain why the power rule holds? Look back to the Handout: *Expanding Powers of Binomials [PDF]* from October 31 (Monday) lecture.
2. $f(x) = 3x^{10}$
3. $f(x) = 3x^{-10}$
4. $f(x) = -\frac{1}{3}x^{-10}$
5. $f(x) = -\frac{1}{x^3}$
6. $f(x) = \frac{42}{x^{10}}$
7. $f(x) = \sqrt[7]{x}$
8. $f(x) = \sqrt[7]{2}$
9. $f(x) = \sqrt[7]{2}x$
10. $f(x) = \frac{5x-2}{x^2+1}$
11. a.) $f(x) = \frac{3-(1/x)}{x+5}$
b.) After that, find the points (x-values) where the graph of the function has a horizontal tangent line.
c.) Use your answer from above to find an equation of the tangent line to the graph $f(x) = \frac{3-(1/x)}{x+5}$ at the point $(-1, 1)$.

3.4 Test 3: Higher derivatives

1. Compute the second derivative $\frac{d^2}{dx^2}(3x^{10} + x^2)$
2. Compute the third derivative $\frac{d^3}{dx^3}(3x^{10} + 5x^2)$

3.5 Test 3: Knowing derivative rules

1. Suppose f is a differentiable function with respect to x . Apply one of the derivative rules and write down the derivative of $5f(x)$.
2. Suppose f is a differentiable function with respect to x . Apply the product rule, and write down the derivative of $x^2 f(x)$.
3. Suppose f is a differentiable function with respect to x . Apply the quotient rule, and write down the derivative of $\frac{f(x)}{2x^5}$.

3.6 Test 3: Figuring out from a graph whether f is continuous, not continuous, and differentiable

1. Sketch an example of a graph that is differentiable everywhere except at $x = 1$ and $x = 8$.
2. Answer True or False. If a graph has a vertical tangent line at $x = 2$, then this graph is differentiable at $x = 2$.
3. Answer True or False. If a graph has a horizontal tangent line at $x = 2$, then this graph is differentiable at $x = 2$.
4. Answer True or False. If f is differentiable at $x = 5$, then f is continuous at $x = 5$.
5. Answer True or False. If f is not continuous at $x = 3$, then f is not differentiable at $x = 3$.
6. Answer True or False. If f is continuous at $x = 2$, then f is differentiable at $x = 2$.
7. Suppose f is a function. What does it mean for f to be differentiable at $x = c$?
8. What does it mean for f to be continuous at $x = c$? Write a precise definition.

3.7 Test 3: Mean Value Theorem Practice

1. Let $f(x)$ be a function that is differentiable on an interval $[2, 5]$, and $f(2) = f(5) = -1$.
 - a. True or false. There is some c between 2 and 5 such that the tangent line to the graph f at $x = c$ is horizontal.
 - b. True or false. There is some c between 2 and 5 such that the tangent line to the graph f at $x = c$ is vertical.
 - c. True or false. There is some c between 2 and 5 such that the tangent line to the graph f at $x = c$ has slope -1 .
2. Suppose you left Saint Peter at 9am and drove to your friend's house 100 miles away. You arrived at 11am, so you completed the trip in two hours.
 - a. What is your *average speed* for the entire trip?
 - b. Sometime between 9 and 11am, did your speedometer ever read exactly 50 mph? In other words, was your *instantaneous speed* ever exactly 50 mph? Yes or no, and why?
3. Suppose f is differentiable on the interval $[1, 7]$. You also know that $f(1) = -3$, $f(6) = 5$, and $f(7) = 15$. Answer True or False. (If false, give a counterexample. If true, explain).
 - a. _____ f must also be continuous on the interval $[1, 7]$.
 - b. _____ There must be at least one number c in $(1, 7)$ where $f'(c) = 3$.
 - c. _____ There must be at least one number c in $(6, 7)$ where $f'(c) = 0$.

3.8 Test 3: Functions and their derivatives

Answer True or False (If false, give a counterexample. If true, explain):

1. T/F: The derivative of a polynomial must be a polynomial.
2. T/F: The derivative of a rational function must be a rational function.
3. T/F: The derivative of an even-degree polynomial must also be an even-degree polynomial.
4. T/F: The derivative of an odd-degree polynomial must also be an odd-degree polynomial.
5. T/F: The derivative of an odd-degree polynomial must be an even-degree polynomial.
6. T/F: The derivative of an even-degree polynomial $f(x)$ (where the degree is 2 or higher) must have at least one zero).
7. T/F: The derivative of an odd-degree polynomial $f(x)$ must have at least one zero.

3.9 From Test 4: Sec 4.4 Composition of functions, Chain Rule (plus all the previous derivative rules from Chapter 4)

1. Write $h(x) = \frac{1}{x+1}$ as a composition of two simpler functions. Then compute $h'(x)$ using Chain Rule.
2. Write $h(x) = \sqrt{3x^2 - x + 1}$ as a composition of two simpler functions. Then compute $h'(x)$ using Chain Rule
3. Use derivative rules to find $f'(x)$. On the test, do not spend a lot of time simplifying once the differentiation steps are completed.
 - (a) $f(x) = (x^2 + 1)^3$. Check your answer with WolframAlpha.
 - (b) $f(x) = \frac{-7}{(2x-3)^2}$. Check your answer with WolframAlpha.
 - (c) $f(x) = \sqrt[3]{(x^2 - 1)^2}$. Check your answer with WolframAlpha.
4. Use your answer from above to find the critical numbers of $f(x) = \sqrt[3]{(x^2 - 1)^2}$. For definition of critical numbers, see Sec 5.1.
5. Read all of Sec 4.4, p284-289. Work through Examples 2-6 (chain rule). Work through Examples 6-9 (how to simplify after differentiation in a clear, readable way).
6. Book Exercises: Sec 4.4 #1-12 (odd numbers first); 47, 59, 60, 61.

3.10 From Test 4: Sec 4.5 Implicit Differentiation

1. Find dy/dx given that $y^3 + y^2 - 5y - x^2 = -4$. (Hint: Differentiate both sides with respect to x , then solve for dy/dx).
2. Find the slope of the tangent line to the graph of $x^2 + 4y^2 = 4$ at the point $(\sqrt{2}, \frac{-1}{\sqrt{2}})$. (Hint: Differentiate both sides with respect to x , then solve for dy/dx , then get the slope you need).

3. Find the equation of the tangent line to the graph of $x^2(x^2 + y^2) = y^2$ at the point $(\sqrt{2}/2, \sqrt{2}/2)$. (Hint: Differentiate both sides with respect to x , then solve for dy/dx , then get the slope you need, then write the equation of the line).
4. Read Sec 4.5 p292-296 for many patiently-worked out implicit diff examples; you'll need to work through them yourself to get fluent with the process.
5. Book Exercises: After you do the reading and work the examples in the section, try Sec 4.5 p297 #1,4, 5, 15, 21.

4 From Test 4

4.1 From Test 4: Sec 4.6 Related Rates

1. Suppose that x and y are both differentiable functions of t , and they are related by the equation $y - x^2 = 3$. Find dy/dt when $x = 1$, given that $dx/dt = 2$ when $x = 1$.
2. I have a 10-ft ladder that is leaning against a wall. At this particular moment, the distance between the bottom of the ladder and the wall is 8 feet. The bottom of the ladder is sliding away from the wall at the rate of 4 ft/second. Hence the top of the ladder is sliding down along the wall. Compute the rate of which the top of the ladder is sliding down at a particular time t . (Hint: See the Falling ladder related rates from Khan Academy).
3. Read Sec 4.6 p300-302. Read the exposition and work through Examples #1-3.
4. Book Exercises: Sec 4.5 #25-31 (odd), 33a, 47; Sec 4.6 #15 (Hint: see Example 2 in the section).
5. Read book Sec 4.6 Ex. 4 (p301).
6. Rework Examples 1-5 (p300-303) on your own (solve it without looking at the book, and then compare to the textbook solution).

4.2 From Test 4: Sec 5.1 Extreme Value Theorem, Extrema

1. Give a complete and accurate statement of the Extreme Value Theorem.
2. Complete the definition. Let f be a function which is defined on c . Then we say that c is a *critical number* of f if
3. True or False: If 5 is a critical number of f , then f has a relative minimum or relative maximum at $x = 5$.
4. True or False: If f has a relative maximum at $x = 2$, then 2 is a critical number of f .
5. Find the absolute minimum and absolute maximum of $f(x) = 3x^4 - 4x^3$ on the interval $[-1, 2]$.
6. Answer True or False.
 - (a) The maximum of a function that is continuous on a closed interval can occur at only one value in the interval.

- (b) If a function is continuous on $[-3, 3]$, then it must have a minimum on the interval.
- (c) If $x = 5$ is a critical number of the polynomial $p(x)$, then $x = 5$ must also be a critical number of the polynomial $f(x) = p(x) + 7$.
- (d) Read Sec 5.1 p314-318. Work through Ex. 2 and 3 on p317-318 (meaning, after you have read through them, copy the problem out and rework it on your own to see if you've really understood and remembered the process).
- (e) Book Exercises: Sec 5.1 (p319) #1-13 - most of these are answered just by looking at a graph, without any calculating. But you'll need to understand the terms and definitions from the reading!
- (f) More Book Exercises: Sec 5.1 p319 #14,16, 17, 19, 21, 45, 46, 47-50 (many of these are graphical exercises that don't require a lot of calculation)

4.3 From Test 4: Sec 5.3 Increase/Decrease/Min/Max


1. Let $f(x) = x^3 - \frac{3}{2}x^2$.
 - (a) What are the critical numbers of f ?
 - (b) Determine the intervals of increase and decrease for f .
 - (c) Classify the critical numbers (relative max/ relative min/ neither).
2. Let $f(x) = 2x^3 - 3x^2 - 36x + 14$.
 - (a) What are the critical numbers of f ?
 - (b) Determine the intervals of increase and decrease for f .
 - (c) Classify the critical numbers (relative max/ relative min/ neither).
3. Find the relative maximum and minimum of

$$f(x) = \frac{x^4 + 1}{x^2}.$$

4. Answer True or False
 - (a) The sum of two increasing functions is increasing.
 - (b) Every n th degree polynomial has $(n - 1)$ critical numbers.
 - (c) There is a relative maximum or minimum at each critical number.
5. Read: Sec 5.3 p329-334.
6. Memorize statements of Theorems 5.5 and 5.6.
7. Work through Examples 1-4 (read, understand, copy the problem, and rework on your own)
8. Book Exercises: Sec 5.3 p335 #1,3,8,11,15,17,19

4.4 From Test 4: Figuring out signs from graphs (Secs. 5.1,5.2,5.4)

The figure shows the graph of a function $f(x)$. Use the graph to determine the sign (positive, negative, or zero) of each of the following.



- | | | |
|------------|-------------|-----------------|
| 1. $f(-4)$ | 6. $f'(-4)$ | 11. $f''(-4)$ |
| 2. $f(-3)$ | 7. $f'(-3)$ | 12. $f''(-3.5)$ |
| 3. $f(0)$ | 8. $f'(0)$ | 13. $f''(0)$ |
| 4. $f(1)$ | 9. $f'(1)$ | 14. $f''(1)$ |
| 5. $f(3)$ | 10. $f'(4)$ | 15. $f''(3)$ |

4.5 From Test 4: Sec 5.4 Concavity

- Complete the definition. Let f be differentiable on an open interval I .
 - The graph of f is *concave up on I* means that
 - The graph of f is *concave down on I* means that
- Give a complete and accurate statement of the test for concavity (Thm 5.7 on p. 339).
- Determine the open intervals on which the graph of $\frac{6}{x^2+3}$ is concave up or down.
- Suppose f is a differentiable function on $(-\infty, \infty)$. Answer True or False.
 - If $f''(x)$ is positive on an interval (a, b) , then $f(x)$ is concave up on (a, b) .
 - If $f''(x)$ is negative on an interval (a, b) , then $f(x)$ is concave down on (a, b) .
 - If $f''(x)$ is positive on an interval (a, b) , then $f(x)$ is increasing on (a, b) .
 - If $f''(x)$ is negative on an interval (a, b) , then $f(x)$ is decreasing on (a, b) .
- If f is continuous everywhere, and f changes from being concave up to concave down at a point where $x = 5$, then this point (where $x = 5$) is a point of inflection of the graph f .

6. Suppose $f(x) = x^N$ where N is an odd positive number larger than 1.
 - (a) Does f have any inflection point/s? How many?
 - (b) Determine the intervals on which the graph of f is concave up or down.
7. Suppose $g(x) = x^N$ where N is an even positive number.
 - (a) Does g have any inflection point/s? How many?
 - (b) Determine the intervals on which the graph of g is concave up or down.
8. Answer True or False. These are exercises # 65-68 from p. 345.
 - (a) Consider a cubic polynomial $h(x) = ax^3 + bx^2 + cx + d$ where $a \neq 0$. This cubic polynomial has precisely one point of inflection.
 - (b) If $f'(c) > 0$, then f is concave upward at $x = c$.
 - (c) If $f''(2) = 0$, then the graph of f must have a point of inflection at $x = 2$.
9. Consider a cubic polynomial $h(x) = ax^3 + bx^2 + cx + d$ where $a \neq 0$. Determine the intervals on which the graph of h is concave up or down.
10. Read Sec 5.4 p338-341. Work the examples and study the figures that go along with them!
11. Book Exercises: Sec 5.4 #1,2,3 (solve these analytically using the second derivative, but look at the graph that's provided to see if your answer matches), #9, 11, 13, 15, 37 and 38, 65.

5 After Test 4

5.1 Optimization problems (Sec 5.7)

1. You want to create an open box (with 4 walls and a base, but no top) with a square base. The surface area has to be 108 square inches. What dimensions (length of the square base, and height) will produce a box with maximum volume?
2. A rectangular page is to contain 24 square inches of print. The margins at the top and bottom of the page are to be $1\frac{1}{2}$ inches, and the margins on the left and right are to be 1 inches. What should be the dimensions of the page be so that the least amount of paper is used? See Sec 5.7 Example 3 p. 366.

5.2 Finding a function given its derivative

1. Could you give me a function whose derivative is equal to 0?
2. Could you give me a function whose derivative is equal to 5?
3. Could you give me a function whose derivative is equal to $2x$?
4. Could you give me a function whose derivative is equal to $3x^2$?