Name : \_\_\_\_\_

(Graded on **correctness** - you can use the internet and share solutions as long as it leads you to learn or re-learn these basic topics - you do not have to show work except on question 11).

## If you can't find it in the book and Googling doesn't help, please ask me!

- 1. The exact value of  $\left(\frac{1}{2} + \frac{1}{3}\right) \div \frac{5}{4}$  is \_\_\_\_\_\_.
- 2. (motivation: partial fraction decomposition) Solve the system of linear equations  $\begin{cases} 2A+B+C &= 1\\ A+2B+C &= 3\\ A+B+2C &= 4 \end{cases}$

A =\_\_\_\_\_\_\_, B =\_\_\_\_\_\_\_, and C =\_\_\_\_\_\_\_

3. (motivation: integral by trig substitution when you need to complete the square) Consider the parabola  $y = x^2 - 3x + 5$ .

The vertex of the parabola is located at y—plane.

4. (motivation: integral by trig substitution when you need to use trig identities) Which of the following equals  $1 - \frac{1}{\tan^2 x + 1}$ ?

A.  $\sin^2 x$ 

B.  $\cos^2 x$ 

C.  $\tan^2 x$ 

D.  $\sec^2 x$ 

(Hint: the back of the book will give you the trig identity  $1 + \tan^2 \theta = \sec^2 \theta$ ).

5. (motivation: polar coordinate, finding area)

The area of a sector with central angle 30°  $(\frac{\pi}{6})$  in a circle of radius 12 m is \_\_\_\_\_ m<sup>2</sup>.

6. For  $-1 \le x \le 1$ , the algebraic expression (without trig expressions) of  $\sin(2\cos^{-1}x)$  is

(Hint: use the identity  $\sin^2(\theta) = 1 - \cos^2(\theta)$ , the definition  $\cos(\cos^{-1} x) = x$ , and the double angle formula  $\sin(2\theta) = 2\sin(\theta)\cos(\theta)$ .

Motivation: See trig substitution method Sec 7.3. Example 1 pg 486-487).

7. (motivation: integrating square of sine/cosine using half-angle formulas) Find the exact values of the following. You can look up half-angle formulas in the back of the book or ask Google.

(a) 
$$\sin^2 \frac{\pi}{8} =$$
 \_\_\_\_\_\_. (b)  $\cos^2 \frac{\pi}{8} =$  \_\_\_\_\_\_.

(b) 
$$\cos^2 \frac{\pi}{8} =$$
 \_\_\_\_\_

8. (motivation: polar coordinate when you need to convert Cartesian coordinate to polar coordinate)

Write the complex number z = 1 - i (that is, the point where x = 1 and y = -1 on the Cartesian plane) in polar form with argument  $\theta$  between 0 and  $2\pi$ :

9. (motivation: reviewing derivative of  $\exp(x)$ ,  $\ln(x)$ ,  $\sin(x)$ ,  $\cos(x)$ , constant, and chain rule, product rule, power rule)

The second derivative of the function  $f(x) = e^{\sin x} + \ln x + \pi^2$  is

$$f''(x) = \underline{\qquad}.$$

10. (motivation: when they want to show that a given sequence is decreasing) Find the interval(s) on which the function  $f(x) = 4x^3 - 15x^2 - 72x + 5$  is decreasing:

- 11. Review u-substitution by doing one of these (or both):
  - watch Khan Academy video https://www.khanacademy.org/math/calculus-home/ap-calculus-ab/ab-antiderivatives-ftc#ab-u-substitution
  - go to Sec 5.5 pages 413-416 and test yourself on Examples 1-6.

Pick two examples from pages 413-416 or the Khan academy video and redo them below without looking at the book/ your note/ video:

(a) \( \int \)

(b) \int \

Calc II Page 3 of 4

12. Evaluate the following.

(a) 
$$\lim_{x\to\infty} \frac{x^2+x-2}{x^2-1} =$$
 \_\_\_\_\_ (compare the coefficients of the highest power of  $x$ ).

(b)  $\lim_{x \to \infty} \frac{\ln x}{\sqrt{x}} =$  \_\_\_\_\_ (practice L'Hospital's Rule).

(c) 
$$\int_0^1 \frac{1}{1+x^2} dx =$$
 \_\_\_\_\_\_\_ (A very common integral. Look this up on the back of the book or ask WolframAlpha).

(d) 
$$\int_0^5 x\sqrt{25-x^2} \, dx = \underline{\qquad}$$
 (review u-substitution).

13. (motivation: used in all the integration skills. It's a good habit to check whether your answer is correct or wrong.)

Tyrion Lannister evaluated the following indefinite integral during the Battle of Calculus :

$$\int x \cos x \, dx = x \sin x + \cos x + C.$$

You don't know how he came up with this result, but you do know whether his answer is correct or not. Why?