

# Sec 9.1 Sequences

A sequence is an ordered collection of objects

## Examples

\* A sequence of letters

\* △ ■ ○ ▲ □ ● △ \_ \_ \_ ...

In Sec 9.1, a sequence is a list of numbers

indexed by the natural numbers 1, 2, 3, 4, ...

Notation:  $\{a_1, a_2, a_3, \dots, a_n, \dots\}$  or  $\{a_n\}$  or  $\{a_n\}_{n=1}^{\infty}$

The index doesn't have to start at 1, ex  $\{a_n\}_{n=0}^{\infty}$  or  $\{a_n\}_{n=5}^{\infty}$

## Examples of sequences

\* 1, 3, 5, 7, 9, ... is the sequence of odd natural numbers

formula  $a_n = 2n - 1$  for  $n = 1, 2, 3, \dots$

\*  $\{a_n\}_{n=1}^{\infty}$  where  $a_n = 2n^2 - 3n + 1$ .

Write the first three terms of  $\{a_n\}_{n=1}^{\infty}$

$$a_1 = 2(1)^2 - 3(1) + 1 = 0$$

$$a_2 = 2(4) - 3(2) + 1 = 3$$

$$a_3 = 2(9) - 3(3) + 1 = 10$$

\* Find a formula for the general term  $a_n$  for the sequence  $\{1, -3, 5, -7, 9, \dots\}$ :

If starting index is  $n=1$ :  $a_n = (2n-1)(-1)^{n+1}$  for  $n=1, 2, 3, \dots$   
 or  $a_n = -(2n-1)(-1)^n$

If starting index is  $n=0$ :  $a_n = (2n+1)(-1)^n$  for  $n=0, 1, 2, \dots$

\* Find a formula for the general term  $a_n$  of the sequence  $\left\{ \frac{3}{5}, -\frac{4}{25}, \frac{5}{125}, -\frac{6}{625}, \frac{7}{3125}, \dots \right\}$ :

• If starting index is  $n=1$ :  $a_1$   $a_2$   $a_3$   $a_4$   $a_5$

• The signs alternate positive & negative, so we need to multiply by  $(-1)^{\text{(something)}}$ .  $a_1$  is positive, so multiply by  $(-1)^{n+1}$  or  $(-1)^{n-1}$ .

• Numerators are  $3, 4, 5, 6, 7, \dots$ :  $(n+2)$  in general  
 $a_1$   $a_2$   $a_3$   $a_4$   $a_5$

• Denominators are  $5, 25, 125, 625, 3125$ :  $5^n$  in general  
 $a_1$   $a_2$   $a_3$   $a_4$   $a_5$

•  $a_n = (-1)^{n+1} \frac{n+2}{5^n}$  for  $n=1, 2, 3, \dots$

• If starting index is  $n=0$ :  $a_n = (-1)^n \frac{n+3}{5^{n+1}}$  or  $\frac{(-1)^n}{5} \frac{n+3}{5^n}$   
 for  $n=0, 1, 2, \dots$

\* The Fibonacci sequence is defined recursively by

$$a_1=1, \quad a_2=1, \quad a_{n+2} = a_n + a_{n+1} \quad \text{for } n=1, 2, 3, \dots$$

each term is the sum of the previous two terms


First few terms of the Fibonacci sequence:

$$\{1, 1, 2, 3, 5, 8, 13, 21, \dots\}$$

\* The sequence  $a_n = \frac{n}{n+1}$  for  $n=1, 2, 3, \dots$

Table:

$n$	1	2	3	4	...	$n$
$a_n$	$\frac{1}{2}$	$\frac{2}{3}$	$\frac{3}{4}$	$\frac{4}{5}$	...	$\frac{n}{n+1}$



(The terms of  $a_n = \frac{n}{n+1}$  seem to approach 1 as  $n$  gets large.)

The difference  $1 - a_n = 1 - \frac{n}{n+1}$

$$= \frac{n+1-n}{n+1}$$
$$= \frac{1}{n+1}$$

Can be made as small as we like by taking large enough  $n$ .

The notation for  $\uparrow$  this is  $\lim_{n \rightarrow \infty} \frac{n}{n+1} = 1$ .

In general, writing  $\lim_{n \rightarrow \infty} a_n = L$  means:

the terms of the sequence  $\{a_n\}$  approach  $L$  as  $n$  becomes large.

New vocab (memorize)

(a number)

\* A sequence  $\{a_n\}$  has limit  $L$  &

we write  $\lim_{n \rightarrow \infty} a_n = L$  or write  $a_n \rightarrow L$  as  $n \rightarrow \infty$

if we can make the terms  $a_n$  as close to  $L$  as

we like by taking  $n$  sufficiently large.

\* If  $\lim_{n \rightarrow \infty} a_n$  exists, say  $\{a_n\}$  converges (or is convergent).  
(is a number)

\* Otherwise, say  $\{a_n\}$  diverges (or is divergent  
or is not convergent).

Ex Is the sequence  $a_n = \frac{2n}{n+1}$  convergent or divergent?

$$\lim_{n \rightarrow \infty} \frac{2n}{n+1} = \lim_{n \rightarrow \infty} \frac{2n}{n+1} \left( \frac{\frac{1}{n}}{\frac{1}{n}} \right)$$

$$= \lim_{n \rightarrow \infty} \frac{2}{\frac{n}{n} + \frac{1}{n}}$$

$$= \lim_{n \rightarrow \infty} \frac{2}{1 + \frac{1}{n}}$$

$$= \frac{(\lim_{n \rightarrow \infty} 2)}$$

$$(\lim_{n \rightarrow \infty} 1) + (\lim_{n \rightarrow \infty} \frac{1}{n})$$

$$= \frac{2}{1+0} = 2$$

(a number)

\* We say:  
 $\{a_n\}$  has limit 2.

\* Since  $\lim_{n \rightarrow \infty} \frac{2n}{n+1}$   
exists, we say  
 $\{a_n\}$  is convergent.

## New vocab

Writing  $\lim_{n \rightarrow \infty} a_n = \infty$  means:

for every positive number  $M$ ,

no matter how big

there is an integer  $N$  such that

if  $n > N$  then  $a_n > M$ .

Say  $\{a_n\}$  diverges to  $\infty$ .

$\lim_{n \rightarrow \infty} a_n = -\infty$  means:

for every positive number  $M$ ,

there is an integer  $N$  such that

if  $n > N$  then  $a_n < -M$ .

Say  $\{a_n\}$  diverges to  $-\infty$ .

Ex Is  $a_n = \frac{-n}{\sqrt{10+n}}$  convergent?

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} \frac{-n}{\sqrt{10+n}} \frac{\left(\frac{1}{n}\right)}{\left(\frac{1}{n}\right)}$$

$$= \lim_{n \rightarrow \infty} \frac{-1}{\sqrt{\frac{10}{n^2} + \frac{n}{n^2}}}$$

$$= \lim_{n \rightarrow \infty} \frac{-1}{\sqrt{\frac{10}{n^2} + \frac{1}{n}}}$$

$$= -\infty$$

numerator =  $-1 \rightarrow -1$  as  $n \rightarrow \infty$

denominator =  $\sqrt{\frac{10}{n^2} + \frac{1}{n}} \rightarrow 0$  as  $n \rightarrow \infty$

\*  $\lim_{n \rightarrow \infty} a_n$  does not exist, so  $\{a_n\}$  diverges (is not convergent).

\*  $\lim_{n \rightarrow \infty} a_n = -\infty$  means  $\{a_n\}$  diverges in a special way:

Say  $\{a_n\}$  diverges to  $-\infty$ .

Thm Let  $f$  be any function.

If  $\lim_{x \rightarrow \infty} f(x) = L$  and  $f(n) = a_n$  when  $n$  is an integer,

then  $\lim_{n \rightarrow \infty} a_n = L$  (Upshot: We can replace  $x$  with  $n$ )

Ex (Application of Thm) Calculate  $\lim_{n \rightarrow \infty} \frac{\ln n}{n}$  :

$$\text{Let } f(x) = \frac{\ln x}{x}.$$

$$\lim_{x \rightarrow \infty} f(x) = \lim_{x \rightarrow \infty} \frac{\ln x}{x} = \lim_{x \rightarrow \infty} \left( \frac{1}{x} \right) \stackrel{(*)}{=} 0$$

↑  
L'Hospital's Rule  
type " $\frac{\infty}{\infty}$ "

Since  $f(n) = a_n$  for  $n = 1, 2, 3, \dots$ , we can apply above Thm:

$$\lim_{n \rightarrow \infty} a_n = \lim_{x \rightarrow \infty} f(x) = 0.$$

↑  
Thm

↑  
by  $(*)$

Thm (Limit laws for convergent sequences)

If  $\{a_n\}$  and  $\{b_n\}$  are convergent sequences and  $c$  is a number,

$$\text{then } * \lim_{n \rightarrow \infty} (a_n + b_n) = \lim_{n \rightarrow \infty} a_n + \lim_{n \rightarrow \infty} b_n$$

$$* \lim_{n \rightarrow \infty} c a_n = c \lim_{n \rightarrow \infty} a_n$$

$$* \lim_{n \rightarrow \infty} (a_n b_n) = \left( \lim_{n \rightarrow \infty} a_n \right) \left( \lim_{n \rightarrow \infty} b_n \right)$$

$$* \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \frac{\lim_{n \rightarrow \infty} a_n}{\lim_{n \rightarrow \infty} b_n} \quad \text{if } \lim_{n \rightarrow \infty} b_n \neq 0$$

## Sandwich Thm (or Squeeze Thm)

If  $a_n \leq b_n \leq c_n$  for  $n \geq N$ , AND

$$* \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} c_n = L$$

THEN  $\lim_{n \rightarrow \infty} b_n = L$

(If  $b_n$  is bounded above & below by two sequences converging to  $L$ , then  $b_n$  converges to  $L$ .)

Thm  
(Special case  
of squeeze Thm)

If  $\lim_{n \rightarrow \infty} |b_n| = 0$  then  $\lim_{n \rightarrow \infty} b_n = 0$ .

EX (of Thm)

Is  $b_n = \frac{(-1)^n}{n}$  convergent?

$$\begin{aligned} \lim_{n \rightarrow \infty} |b_n| &= \lim_{n \rightarrow \infty} \left| \frac{(-1)^n}{n} \right| \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \\ &= 0 \end{aligned}$$

By Squeeze Thm,  $\lim_{n \rightarrow \infty} b_n = 0$ . So  $\lim_{n \rightarrow \infty} b_n$  exists.

So  $\{b_n\}$  is convergent.

Thm If  $\lim_{n \rightarrow \infty} a_n = L$  and function  $f$  is continuous at  $L$ ,

$$\text{then } \lim_{n \rightarrow \infty} f(a_n) = f\left(\lim_{n \rightarrow \infty} a_n\right) = f(L)$$

Upshot: can bring  $\lim_{n \rightarrow \infty}$  inside brackets if  $f$  is continuous at  $L$ .

Ex (of thm)

$$\lim_{n \rightarrow \infty} \sin\left(\frac{\pi}{n}\right) = ?$$

$$\text{Let } a_n := \frac{\pi}{n}. \text{ Then } \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} \frac{\pi}{n} = 0.$$

Let  $f(x) := \sin x$ . Then  $f(x)$  is continuous at  $0$ .

$$\text{So } \lim_{n \rightarrow \infty} f(a_n) = \underset{\substack{\uparrow \\ \text{by Thm}}}{f}\left(\lim_{n \rightarrow \infty} a_n\right) = f(0)$$

$$\lim_{n \rightarrow \infty} \sin\left(\frac{\pi}{n}\right) = \underset{\substack{\uparrow \\ \text{by Thm}}}{\sin}\left(\lim_{n \rightarrow \infty} \frac{\pi}{n}\right) = \sin(0) = 0.$$



## New vocab

$a_n = r^n$  (like  $a_n = (\frac{1}{2})^n$ ,  $a_n = (-2)^n$ ,  $a_n = 1^n$ ,  $a_n = (-1)^n$ )  
"ratio"  
is called a geometric sequence.

Ex \*  $\lim_{n \rightarrow \infty} (\frac{1}{2})^n = 0$ , say  $\{\frac{1}{2^n}\}$  converges to 0

\*  $\lim_{n \rightarrow \infty} 1^n = 1$ , say  $\{1\}$  converges to 1

\*  $\lim_{n \rightarrow \infty} 2^n = \infty$  <sup>not a number</sup>, say  $\{2^n\}$  diverges to  $\infty$   
does not exist

\*  $\lim_{n \rightarrow \infty} (\frac{-2}{3})^n = 0$  by Sandwich (or squeeze) Thm.  
Say  $\{(\frac{-2}{3})^n\}$  converges to 0.

\*  $\lim_{n \rightarrow \infty} (-1)^n$  doesn't exist. Say  $\{(-1)^n\}$  diverges.

\*  $\lim_{n \rightarrow \infty} (\frac{-3}{2})^n$  doesn't exist. Say  $\{(\frac{-3}{2})^n\}$  diverges.

Fact The geometric sequence  $\{r^n\}$

is convergent if  $-1 < r \leq 1$ : (like  $r = 1, \frac{1}{2}, -\frac{2}{3}$ )

$$\lim_{n \rightarrow \infty} r^n = 0 \quad \text{if } -1 < r < 1$$

$$\lim_{n \rightarrow \infty} 1^n = 1$$

$\{r^n\}$  diverges if  $r \leq -1$  or  $1 < r$  (like  $r = -1, \frac{3}{2}, 2$ )

New vocab

\*  $\{a_n\}$  is increasing if  $a_n < a_{n+1}$  for all  $n \geq 1$ :

$$a_1 < a_2 < a_3 < \dots$$

\*  $\{a_n\}$  is decreasing if  $a_n > a_{n+1}$  for all  $n \geq 1$ :

$$a_1 > a_2 > a_3 > \dots$$

\*  $\{a_n\}$  is monotonic if it is either increasing or decreasing.

Ex Is  $\frac{3}{n+5}$  monotonic?

$$a_1 = \frac{3}{6} > a_2 = \frac{3}{7} > a_3 = \frac{3}{8} > \dots$$

$$a_n = \frac{3}{n+5} > a_{n+1} = \frac{3}{n+6} \text{ for all } n=1, 2, \dots$$

So  $\{a_n\}$  is decreasing, so  $\{a_n\}$  is monotonic.

New vocab

\*  $\{a_n\}$  is bounded above if there is a number  $M$

such that  $a_n \leq M$  for all  $n \geq 1$ .

\*  $\{a_n\}$  is bounded below if there is a number  $m$

such that  $m \leq a_n$  for all  $n \geq 1$ .

\* Say  $\{a_n\}$  is bounded if

$\{a_n\}$  is bounded above and below.

Ex  $\frac{3}{n+5}$  lower bounds: upper bounds:  
 $0, -\frac{1}{2}$   $\frac{3}{6}, 1, 1000$

## Monotonic Sequence Thm

If  $\{a_n\}$  is bounded and monotonic, then  $\{a_n\}$  converges.

Ex  $\left\{\frac{3}{n+5}\right\}$  is decreasing and bounded,

so by the monotonic sequence thm,

$\left\{\frac{3}{n+5}\right\}$  converges.

True or false?

1. If a sequence  $\{a_n\}$  is bounded, then  $\{a_n\}$  is convergent.

False. Counterexample:

$\{1, -1, 1, -1, \dots\}$  is bounded by  $-1$  and  $1$  but it diverges.

2. If  $\{a_n\}$  is monotonic, then  $\{a_n\}$  is convergent.

False. Counterexample:

Let  $a_n = n$ . Then  $\{a_n\}$  is increasing but  $\lim_{n \rightarrow \infty} a_n = \infty$

so  $\{a_n\}$  is divergent.

3. If  $\{a_n\}$  is convergent, then  $\{a_n\}$  is monotonic.

False. Counterexample:  $a_n = \frac{(-1)^n}{n}$  is convergent but not monotonic (neither increasing nor decreasing).