

Recall (Sec 6.3 Thm 9) "The best approximation theorem"

Let W be a subspace of $\mathbb{R}^n$, let $\vec{b}$ be a vector in $\mathbb{R}^n$.

Then the closest point in W to $\vec{b}$ is

$\hat{\vec{b}} = \text{proj}_W \vec{b}$, the orthogonal projection of $\vec{b}$ onto W ,

i.e. $\|\vec{b} - \hat{\vec{b}}\| < \|\vec{b} - \vec{v}\|$ for any $\vec{v}$ in W that's not equal to $\hat{\vec{b}}$.

Sec 6.5 Least-squares problems

Motivation:

- For applications, we often work with systems

$$A\vec{x} = \vec{b}$$

that have no solutions (i.e. are inconsistent)

- We can approximate a solution by finding an $\vec{x}$

that makes $A\vec{x}$ as close as possible to $\vec{b}$,

i.e. by finding an $\vec{x}$ that makes

$$\|\vec{b} - A\vec{x}\| \text{ as small as possible}$$

This process is called the general least-squares problem

- "least-squares" comes from the fact that

$$\|\vec{b} - A\vec{x}\| = \sqrt{\text{sum of squares}}$$

How to solve the general least-squares problem?

Let A be an $m \times n$ matrix and $\vec{b}$ in $\mathbb{R}^m$


Consider the system $A \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix}$

Let W denote $\text{Col } A = \text{Span} \{ \text{columns of } A \}$
 (see Sec 4.2) $\Rightarrow \{ \vec{v} : A\vec{x} = \vec{v} \text{ for some } \vec{x} \text{ in } \mathbb{R}^n \}$

Ex: $A = \begin{bmatrix} 4 & 0 \\ 0 & 2 \\ 1 & 1 \end{bmatrix} \quad \vec{b} = \begin{bmatrix} 2 \\ 0 \\ 11 \end{bmatrix}$
 (Example 1)

Note $\left[\begin{array}{cc|c} 4 & 0 & 2 \\ 0 & 2 & 0 \\ 1 & 1 & 11 \end{array} \right] \rightarrow \left[\begin{array}{cc|c} 4 & 0 & 2 \\ 0 & 2 & 0 \\ -4 & -4 & -44 \end{array} \right] \rightarrow \left[\begin{array}{cc|c} 4 & 0 & 2 \\ 0 & 4 & 0 \\ 0 & -4 & -42 \end{array} \right] \rightarrow \left[\begin{array}{cc|c} 4 & 0 & 2 \\ 0 & 4 & 0 \\ 0 & 0 & -42 \end{array} \right]$ so $A \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ 0 \\ 11 \end{bmatrix}$ is inconsistent

Let $\hat{\vec{b}}$ denote $\text{proj}_W \vec{b}$, the orthogonal projection of $\vec{b}$ onto W .



Q: Is $\hat{\vec{b}}$ in W ?

Ans: Yes, by def

So $\hat{\vec{b}}$ is in $\text{Col } A$, so there is $\hat{\vec{x}}$ in $\mathbb{R}^n$ such that $A\hat{\vec{x}} = \hat{\vec{b}}$.

Since $\hat{\vec{b}} = \text{proj}_W \vec{b}$ is the closest point in $W = \text{Col } A$ to $\vec{b}$,

$A\hat{\vec{x}} = \hat{\vec{b}}$ is the closest point in $W = \text{Col } A$ to $\vec{b}$.

That is, $\| \vec{b} - \hat{\vec{b}} \| \leq \| \vec{b} - \vec{v} \|$ for all $\vec{v}$ in $W = \text{Col } A$

i.e. $\| \vec{b} - A\hat{\vec{x}} \| \leq \| \vec{b} - A\vec{x} \|$ for all $\vec{x}$ in $\mathbb{R}^n$


Def

If A is $m \times n$ and $\mathbf{b}$ is in $\mathbb{R}^m$, a **least-squares solution** of $A\mathbf{x} = \mathbf{b}$ is an $\hat{\mathbf{x}}$ in $\mathbb{R}^n$ such that

$$\|\mathbf{b} - A\hat{\mathbf{x}}\| \leq \|\mathbf{b} - A\mathbf{x}\|$$

for all $\mathbf{x}$ in $\mathbb{R}^n$.

Note: $\hat{\mathbf{x}}$ is a least-squares solution of $A\vec{x} = \vec{b}$ iff $A\hat{\mathbf{x}} = \hat{\mathbf{b}}$



How do we solve $A\hat{\mathbf{x}} = \hat{\mathbf{b}}$?

Suppose $\hat{\mathbf{x}}$ is a vector in $\mathbb{R}^n$ satisfying $A\hat{\mathbf{x}} = \hat{\mathbf{b}}$.

By the Orthogonal Decomposition Theorem (Sec 6.3),

$$\vec{z} = \vec{b} - \hat{\mathbf{b}} \text{ is orthogonal to } W = \text{Col } A = \text{Span}\{\text{cols of } A\} \\ = \vec{b} - A\hat{\mathbf{x}}$$

so $\vec{b} - A\hat{\mathbf{x}}$ is orthogonal to each column of A .

So... If $\vec{a}$ in $\mathbb{R}^m$ is any column of A ,

$$\text{then } 0 = \vec{a} \cdot (\vec{b} - A\hat{\mathbf{x}}) = \vec{a}^T (\vec{b} - A\hat{\mathbf{x}}) \quad \text{def of inner product}$$

$$\text{Ex: } \begin{bmatrix} 4 & 0 & 1 \end{bmatrix} (\vec{b} - A\hat{\mathbf{x}}) = 0$$

Since $\vec{a}^T$ is a row of A^T , we have ...

$$\begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix} = A^T (\vec{b} - A\hat{\mathbf{x}}) \quad \text{Ex: } \begin{bmatrix} 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 4 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix} (\vec{b} - A\hat{\mathbf{x}})$$

$$\text{Thus } \vec{0} = A^T \vec{b} - A^T A \hat{\mathbf{x}}, \text{ i.e. } A^T A \hat{\mathbf{x}} = A^T \vec{b}.$$

Def The matrix equation $A^T A \hat{x}^{(*)} = A^T \vec{b}$ represents

a system of equations called the normal equations for $A\vec{x} = \vec{b}$

Ex: The normal equations for our $A\vec{x} = \vec{b}$ are

$$A^T A \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = A^T \vec{b}$$

$$\begin{bmatrix} 4 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} 4 & 0 \\ 0 & 2 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 4 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 0 \\ 11 \end{bmatrix}$$

$$\begin{bmatrix} 17 & 1 \\ 1 & 5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 19 \\ 11 \end{bmatrix}$$

Thm The set of least-squares solutions of $A\vec{x} = \vec{b}$
(Thm 13) is equal to the (nonempty) set
 $\left\{ \text{solutions of the normal equations } A^T A \vec{x} = A^T \vec{b} \right\}$

$$\text{Ex} \quad \left[\begin{array}{cc|c} 17 & 1 & 19 \\ 1 & 5 & 11 \end{array} \right] \rightsquigarrow \left[\begin{array}{cc|c} 1 & 0 & 1 \\ 0 & 1 & 2 \end{array} \right]$$

For this example of A and $\vec{b}$, the equation $A\vec{x} = \vec{b}$
has a unique least-squares solution, $\hat{x} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$

EXAMPLE 2 Find a least-squares solution of $A\mathbf{x} = \mathbf{b}$ for

$$A = \begin{bmatrix} 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} -3 \\ -1 \\ 0 \\ 2 \\ 5 \\ 1 \end{bmatrix}$$

SOLUTION Compute

$$A^T A = \begin{bmatrix} 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 6 & 2 & 2 & 2 \\ 2 & 2 & 0 & 0 \\ 2 & 0 & 2 & 0 \\ 2 & 0 & 0 & 2 \end{bmatrix}$$

$$A^T \mathbf{b} = \begin{bmatrix} 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} -3 \\ -1 \\ 0 \\ 2 \\ 5 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 \\ -4 \\ 2 \\ 6 \end{bmatrix}$$

The augmented matrix for $A^T A \mathbf{x} = A^T \mathbf{b}$ is

$$\begin{bmatrix} 6 & 2 & 2 & 2 & 4 \\ 2 & 2 & 0 & 0 & -4 \\ 2 & 0 & 2 & 0 & 2 \\ 2 & 0 & 0 & 2 & 6 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 0 & 1 & 3 \\ 0 & 1 & 0 & -1 & -5 \\ 0 & 0 & 1 & -1 & -2 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

The general solution is $x_1 = 3 - x_4$, $x_2 = -5 + x_4$, $x_3 = -2 + x_4$, and x_4 is free. So the general least-squares solution of $A\mathbf{x} = \mathbf{b}$ has the form

$$\hat{\mathbf{x}} = \begin{bmatrix} 3 \\ -5 \\ -2 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} -1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$$

The next theorem gives useful criteria for determining when there is only one least-squares solution of $A\mathbf{x} = \mathbf{b}$. (Of course, the orthogonal projection $\hat{\mathbf{b}}$ is always unique.)

Let A be an $m \times n$ matrix. The following statements are logically equivalent:

- The equation $A\mathbf{x} = \mathbf{b}$ has a unique least-squares solution for each $\mathbf{b}$ in $\mathbb{R}^m$.
- The columns of A are linearly independent.
- The matrix $A^T A$ is invertible.

When these statements are true, the least-squares solution $\hat{\mathbf{x}}$ is given by

$$\hat{\mathbf{x}} = (A^T A)^{-1} A^T \mathbf{b} \quad (4)$$

Ex 1 has a unique least-squares solution since $A^T A$ is invertible
Ex 2 does not since $A^T A$ is not invertible.

When a least-squares solution $\hat{\mathbf{x}}$ is used to produce $A\hat{\mathbf{x}}$ as an approximation to $\mathbf{b}$, the distance from $\mathbf{b}$ to $A\hat{\mathbf{x}}$ is called the **least-squares error** of this approximation.

EXAMPLE 3 Given A and $\mathbf{b}$ as in Example 1, determine the least-squares error in the least-squares solution of $A\mathbf{x} = \mathbf{b}$.




FIGURE 3

SOLUTION From Example 1,

$$\mathbf{b} = \begin{bmatrix} 2 \\ 0 \\ 11 \end{bmatrix} \quad \text{and} \quad A\hat{\mathbf{x}} = \begin{bmatrix} 4 & 0 \\ 0 & 2 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 4 \\ 4 \\ 3 \end{bmatrix}$$

Hence

$$\mathbf{b} - A\hat{\mathbf{x}} = \begin{bmatrix} 2 \\ 0 \\ 11 \end{bmatrix} - \begin{bmatrix} 4 \\ 4 \\ 3 \end{bmatrix} = \begin{bmatrix} -2 \\ -4 \\ 8 \end{bmatrix}$$

and

$$\|\mathbf{b} - A\hat{\mathbf{x}}\| = \sqrt{(-2)^2 + (-4)^2 + 8^2} = \sqrt{84}$$

The least-squares error is $\sqrt{84}$. For any $\mathbf{x}$ in $\mathbb{R}^2$, the distance between $\mathbf{b}$ and the vector $A\mathbf{x}$ is at least $\sqrt{84}$. See Figure 3. Note that the least-squares solution $\hat{\mathbf{x}}$ itself does not appear in the figure. ■

— end of PDF —