

Read pg 759-760:

Suppose for $|x-a| < R$, we have

$$f(x) = c_0 + c_1(x-a) + c_2(x-a)^2 + c_3(x-a)^3 + c_4(x-a)^4 + \dots,$$

then $f(a) = c_0$. We can differentiate both sides with respect to x to get

$$f'(x) = c_1 + 2c_2(x-a) + 3c_3(x-a)^2 + 4c_4(x-a)^3 + \dots,$$

then $f'(a) = c_1$. Again, we have

$$f''(x) = 2c_2 + 2 \times 3c_3(x-a) + 3 \times 4c_4(x-a)^2 + \dots,$$

then $f''(a) = 2c_2$. Apply the procedure one more time to obtain

$$f'''(x) = 2 \times 3c_3 + 2 \times 3 \times 4c_4(x-a) + \dots,$$

then $f'''(a) = 2 \times 3c_3$. By now you can see the pattern. If we continue to differentiate and substitute $x = a$, we obtain

$$f^{(n)}(a) = n! c_n.$$

That is

$$c_n = \frac{f^{(n)}(a)}{n!}.$$

Example:

If $f(x) = \sum_{n=0}^{\infty} c_n(x-5)^n$ for all x , write a formula for c_8 . Answer: follow Theorem 5 (pg 760).

Theorem (Sec 11.10 page 760)

If f has a power series representation at $x = a$, that is, if

$$f(x) = \sum_{n=0}^{\infty} c_n (x-a)^n \quad \text{for} \quad |x-a| < R,$$

then its coefficients are given by

$$c_n = \frac{f^{(n)}(a)}{n!}.$$

$$\begin{aligned} f(x) &= \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x-a)^n \\ &= f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \frac{f'''(a)}{3!}(x-a)^3 + \dots \end{aligned}$$

The series is called the **Taylor series** of the function f at $x = a$. For the special case when $a = 0$, the Taylor series becomes

$$f(x) = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \dots$$

This case arises frequently enough that it is given the special name **Maclaurin series**.

Example (check solution of Example 1, pg 760):

Find the **Maclaurin series** of the function $f(x) = e^x$ and its interval of convergence.

Recall the proof that e is irrational (from Sec 11.5, Alternating Series Estimate Theorem):

We needed the fact that $1/e$ is equal to the series _____ ?

Consider the partial sums

$$T_n(x) = \sum_{k=0}^n \frac{f^{(k)}(a)}{k!} (x-a)^k$$

Notice that T_n is a polynomial of degree n called the n th-degree **Taylor polynomial** of f at $x = a$.

Example:

Consider the function $f(x) = \ln x$.

- Find the **Taylor polynomial of order 2** centered at $x = 1$ for $f(x) = \ln x$.
- Estimate the value of $\ln 1.05$. Give an upper bound of the error of this approximation.