

A **sequence** is an ordered collection of objects.

Example: A sequence of musical notes

Example: You may have seen a pattern recognition “IQ” puzzle similar to this:



In calculus, a **sequence** can be thought of as a list of numbers indexed by the natural numbers 1,2,3,4, ... :

$$a_1, a_2, a_3, a_4, \dots, a_n, \dots$$

or a function from the natural numbers to the real numbers.

The number a_1 is called the first term, a_2 is the second term, and in general a_n is the n th term.

The sequence $\{a_1, a_2, a_3, \dots\}$ is also denoted by

$$\{a_n\} \text{ or } \{a_n\}_{n=1}^{\infty}$$

Note that n doesn't have to start at 1.

Example: 1, 3, 5, 7, 9, ... is a sequence of odd natural numbers.

A sequence can be finite or infinite, but for this class we're mostly interested in infinite sequences and what happens as you look further down the sequence.

Some sequences can be defined by giving an **explicit formula** for the n th term. For example, the **Geometric Sequence** (See Sec 11.1 Example 11):

$$a_n = a_1 r^{n-1}.$$

Each subsequent number is determined by multiplying the previous term by a fixed, nonzero number.

Task 1 Example:

Write the first four terms of the sequence $\{a_n\}_{n=1}^{\infty}$ where $a_n = 2n^2 - 3n + 1$.

Task 2 Example: Find a formula for the general term a_n for the sequence of odd natural numbers. Then find a formula for the sequence $\{1, -3, 5, -7, 9, \dots\}$.

Example: Find a formula for the general term a_n of the sequence

$$\left\{ \frac{3}{5}, -\frac{4}{25}, \frac{5}{125}, -\frac{6}{625}, \frac{7}{3125}, \dots \right\}$$

assuming that the pattern of the first few terms continues.

Example: 1,1,2,3,5,8,13, ... the **Fibonacci Sequence** can be defined by:

$$a_1 = 1, a_2 = 1 \text{ and } a_{n+2} = a_n + a_{n+1} \text{ for all } n \geq 1.$$

Each subsequent number is the sum of the previous two.

Task 3 Example: Consider the sequence $a_n = \frac{n}{n+1}$. (Create a table with Desmos.com)

It appears that the terms of the sequence $a_n = \frac{n}{n+1}$ are approaching 1 as n becomes large. In fact, the difference

$$1 - \frac{n}{n+1} = \frac{1}{n+1}$$

can be made as small as we like by taking n sufficiently large. We indicate this by writing

$$\lim_{n \rightarrow \infty} \frac{n}{n+1} = 1$$

In general, the notation

$$\lim_{n \rightarrow \infty} a_n = L$$

means that the terms of the sequence $\{a_n\}$ approach L as n becomes large.

Task 4 Definition (copy from page 696)

A sequence $\{a_n\}$ has the limit L and we write

if we can make the terms a_n as close to L as we like by taking n sufficiently large. If $\lim_{n \rightarrow \infty} a_n$ exists, we say the sequence **converges** (or is **convergent**). Otherwise, we say the sequence **diverges** (or is **divergent**).

Task 5: Determine whether the sequence $\{a_n\}$ where $a_n = \frac{n}{\sqrt{10+n}}$ is convergent. (Ex.4 pg698)

Task 6 Theorem (from book, Theorem 3 on pg 697)

If $\lim_{x \rightarrow \infty} f(x) = L$ and $f(n) = a_n$ when n is an integer, then _____ .

Why?

Task 7 Ex: Determine whether the sequence $\{a_n\}$ where $a_n = \frac{\ln n}{n}$ is convergent. (Ex. 6 pg 698)

Theorem

If $\{a_n\}$ and $\{b_n\}$ are convergent sequences and c is a constant, then

- $\lim_{n \rightarrow \infty} (a_n + b_n) = \lim_{n \rightarrow \infty} a_n + \lim_{n \rightarrow \infty} b_n$.
- $\lim_{n \rightarrow \infty} ca_n = c \lim_{n \rightarrow \infty} a_n$.
- $\lim_{n \rightarrow \infty} (a_n b_n) = \lim_{n \rightarrow \infty} a_n \cdot \lim_{n \rightarrow \infty} b_n$.
- $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \frac{\lim_{n \rightarrow \infty} a_n}{\lim_{n \rightarrow \infty} b_n}$ provided that $\lim_{n \rightarrow \infty} b_n \neq 0$.

Theorem (Squeeze Theorem for sequences)

If $a_n \leq b_n \leq c_n$ for $n \geq N$ and $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} c_n = L$, then $\lim_{n \rightarrow \infty} b_n = L$.

Task 8: Determine whether the sequence $\{a_n\}$ where $a_n = \frac{(-1)^n}{n}$ is convergent. (Ex. 8 pg699).

Task 9 Theorem (read carefully, from book, Theorem 7 on page 699)

If $\lim_{n \rightarrow \infty} a_n = L$ and the function f is continuous at L , then

$$\lim_{n \rightarrow \infty} f(a_n) = f\left(\lim_{n \rightarrow \infty} a_n\right) = f(L).$$

Task 10: Determine whether the sequence $\{a_n\}$ where $a_n = \sin\left(\frac{\pi}{n}\right)$ is convergent. (Ex. 9 pg 699).

Task 11: Re-read Thm 3 and Thm 7 from the book. What are the differences between the two?