

Part 2 NAME: \_\_\_\_\_

Notes


**Recall**

The **improper integral** is used for cases in which

- The **interval** of integration is **infinite** or
- The **integrand** has an **infinite discontinuity** on the interval of integration.

**Infinite Discontinuity**

Consider the integral  $\int_c^1 \frac{1}{\sqrt{x}} dx$ , where  $0 < c < 1$ .



$$\int_c^1 \frac{1}{\sqrt{x}} dx = (2\sqrt{x})_c^1 = 2 - 2\sqrt{c}$$

$c = \frac{1}{4}$	$c = \frac{1}{9}$	$c = \frac{1}{16}$		$c \rightarrow 0^+$ or
$2 - 2\sqrt{\frac{1}{4}}$	$2 - 2\sqrt{\frac{1}{9}}$	$2 - 2\sqrt{\frac{1}{16}}$		or

We express this result as

$$\int_0^1 \frac{1}{\sqrt{x}} dx = 2$$

which is an improper integral because 0 leads to a zero-denominator.

**Definition Type 2: Improper Integrals with an Unbounded Integrand (copy from pg 531)**

1. If  $f$  is continuous on  $[a, b)$  and is discontinuous at  $b$ , then

$$\int_a^b f(x) dx = \underline{\hspace{10cm}}$$

or  $\underline{\hspace{10cm}}$ ,

provided this limit exists (as a finite number).

2. If  $f$  is continuous on  $(a, b]$  and is discontinuous at  $a$ , then

$$\int_a^b f(x) dx = \underline{\hspace{10cm}},$$

or  $\underline{\hspace{10cm}}$ ,

provided this limit exists (as a finite number).

The improper integrals  $\int_a^b f(x) dx$  is called

- **convergent** if the corresponding limit exists and
- **divergent** if the limit does not exist.

3. If  $f$  has a discontinuity at  $c$ , where  $a < c < b$ , and both  $\int_a^c f(x) dx$  and  $\int_c^b f(x) dx$  are convergent, then we define

$$\int_a^b f(x) dx = \underline{\hspace{10cm}}.$$

Example: Sketch and evaluate  $\int_0^3 \frac{1}{\sqrt{9-x^2}} dx$ . (Hint: first evaluate the indefinite integral using trig substitution)

Example (copy Example 8 pg 532: Use integration by parts and l'Hopital's Rule):

a) Sketch and evaluate  $\int_0^1 \ln x \, dx$ . (Explain your work involving L'Hospital.)

b) Evaluate  $\lim_{t \rightarrow 0^+} t (\ln t)^2$  or  $\lim_{t \rightarrow 0^+} t (\ln t)^3$  (Apply L'Hospital multiple times.)

c) Evaluate  $\int_0^1 (\ln x)^2 \, dx$  or  $\int_0^1 (\ln x)^3 \, dx$ . Use the previous part (b) to help you get your answer.


d) Evaluate  $\int_0^1 (\ln x)^n \, dx$  for  $n = 2, 3, 4, 5, 6$  using a computing tool (like WolframAlpha).  
Predict the value for any  $n$ .

A Comparison Test for Improper Integrals

**Theorem Comparison Theorem (copy from pg 533)**

Suppose that  $f$  and  $g$  are continuous functions with  $f(x) \geq g(x) \geq 0$  for  $x \geq a$ .

1. If  $\int_a^\infty f(x) dx$  is convergent, then  $\int_a^\infty g(x) dx$  is \_\_\_\_\_.
2. If  $\int_a^\infty g(x) dx$  is divergent, then  $\int_a^\infty f(x) dx$  is \_\_\_\_\_.



**Caution**

- If  $\int_a^\infty g(x) dx$  is convergent, then \_\_\_\_\_.
- If  $\int_a^\infty f(x) dx$  is divergent, then \_\_\_\_\_.

Example 10 (copy from pg 534):

Use the comparison test to show that the integral  $\int_1^\infty \frac{1+e^{-x}}{x} dx$  is convergent/divergent.

**Predict a Limit Comparison Test for Improper Integrals!**

(Hint: the Limit Comparison Test for series from Sec 11.4).