

17.6 Surface integrals

I. Parametrized surfaces

* A parametrization of a curve C in \mathbb{R}^2 :

$$\vec{r}(t) = \langle x(t), y(t) \rangle \text{ for } a \leq t \leq b.$$

one parameter t

two dependent variables

* A parametrization of a surface S in \mathbb{R}^3 :


$$\vec{r}(u, v) = \langle x(u, v), y(u, v), z(u, v) \rangle$$

Two parameters u, v


Three dependent variables

where the parameters u and v vary over a rectangle

$$R = \{(u, v) : a \leq u \leq b, c \leq v \leq d\}$$



each point (u, v) in R
is sent to a point
 $\vec{r}(u, v)$ on S



(Note:
parametric
description of
a surface is
not unique,
just like
parametric
description of
a curve is
not unique)




Examples of parametrized surfaces

$$1. \text{ Cylinder } S \quad \{(x, y, z) : x = 5 \cos \theta, y = 5 \sin \theta, 0 \leq \theta \leq \pi, 2 \leq z \leq 6\}$$

(MML #1) half a cylinder, 1st and 2nd quadrant only

A possible parametrization of S is to let $u = \theta$, $v = z$

$$\begin{aligned}\vec{r}(u, v) &= \langle x(u, v), y(u, v), z(u, v) \rangle \\ &= \langle 5 \cos u, 5 \sin u, v \rangle \\ \text{for } 0 &\leq u \leq \pi \text{ and } 2 \leq v \leq 6\end{aligned}$$



$$2. \text{ Plane } S \quad \left\{ (x, y, z) : \underbrace{3x - 2y + z = 2} \right\}$$


Let $u = x$, $v = y$, and $\underbrace{z = -3x + 2y + 2}_{z = -3u + 2v + 2}$
 x and y can be
any number

A possible parametrization of S is


$$\vec{r}(u, v) = \left\langle u, v, \underbrace{-3u + 2v + 2}_{z} \right\rangle \text{ for } -\infty < u < \infty \text{ and } -\infty < v < \infty.$$

II. Surface integral of a scalar-valued function $\iint_S f(x, y, z) \, dS$

Idea: partition rectangle $R = \{(u, v) : a \leq u \leq b, c \leq v \leq d\}$ into small rectangles



$$\text{partition rectangle } R = \{(u, v) : a \leq u \leq b, c \leq v \leq d\}$$



Each small rectangle gets

sent by $\vec{r}(u, v) = \langle x(u, v), y(u, v), z(u, v) \rangle$


to a curved patch S_k on the surface S .

Let $P(a, b)$ be the lower left corner point of the small rectangle.

$\vec{r}(u, v)$ sends $P(a, b)$ to a point $\underbrace{P(x(a, b), y(a, b), z(a, b))}_{\text{on } S_k}$ on S_k .

Take the sum of $f(\vec{r})$ over all lower left corners of the small rectangles R_k multiplied by area of S_k

$$\sum_{\text{over all small rectangles}} f(x(a_k, b_k), y(a_k, b_k), z(a_k, b_k)) (\text{area of } S_k)$$



$$\vec{t}_v = \frac{\partial \vec{r}}{\partial v} = \left\langle \frac{\partial x}{\partial v}, \frac{\partial y}{\partial v}, \frac{\partial z}{\partial v} \right\rangle$$

tangent vector corresponding to change in v (keeping u constant)

$$\vec{t}_u = \frac{\partial \vec{r}}{\partial u} = \left\langle \frac{\partial x}{\partial u}, \frac{\partial y}{\partial u}, \frac{\partial z}{\partial u} \right\rangle$$

tangent vector corresponding to change in u (keeping v constant)

this parallelogram in the tangent plane

$$\begin{aligned} \text{area} &= |\Delta u \vec{t}_u \times \Delta v \vec{t}_v| \\ &= |\vec{t}_u \times \vec{t}_v| \Delta u \Delta v \end{aligned}$$

Def The surface integral of $f(x, y, z)$ on surface S , denoted $\iint_S f(x, y, z) dS$ is the limit of the sum as the sizes of the small rectangles go to 0.

To compute surface integral, we first convert it to a double integral over the rectangle R :

(i) using a parametric description of surface S :

$$\iint_S f(x, y, z) dS = \iint_R f(x(u, v), y(u, v), z(u, v)) \left| \vec{t}_u \times \vec{t}_v \right| dA$$

(ii) If S is explicitly defined by $z = g(x, y)$ for (x, y) in region R , then $\vec{t}_u = \vec{t}_x = \langle 1, 0, z_x \rangle$ $\vec{t}_v = \vec{t}_y = \langle 0, 1, z_y \rangle$

$$\vec{t}_x \times \vec{t}_y = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 0 & z_x \\ 0 & 1 & z_y \end{vmatrix} = \langle -z_x, -z_y, 1 \rangle$$

$$\left| \vec{t}_x \times \vec{t}_y \right| = \sqrt{z_x^2 + z_y^2 + 1}$$

So $\iint_S f dS = \iint_R f(x, y, g(x, y)) \sqrt{z_x^2 + z_y^2 + 1} dA$

Both are the usual 2D double integral from sec 16.2-16.3.

If $f(x, y, z) = 1$, then $\iint_S f(x, y, z) dS = \text{surface area of } S$

Ex 2(a) and MML # 3 .



Find the area of the surface of the half cylinder

$$\{(r, \theta, z) : r=7, 0 \leq \theta \leq \pi, 0 \leq z \leq 8\}$$

using a parametric description of S.

So 1: Parametrize S:


$$\begin{aligned}\vec{r}(u, v) &= \langle x(u, v), y(u, v), z(u, v) \rangle \\ &= \langle 7 \cos u, 7 \sin u, v \rangle \\ \text{for } 0 \leq u &\leq \pi \text{ and } 0 \leq v \leq 8\end{aligned}$$



$$\text{Surface Area} = \iint_S 1 dS = \iint_R 1 |\vec{t}_u \times \vec{t}_v| dA$$

$$\vec{t}_u \times \vec{t}_v = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ x_u & y_u & z_u \\ x_v & y_v & z_v \end{vmatrix} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -7 \sin u & 7 \cos u & 0 \\ 0 & 0 & 1 \end{vmatrix} = \hat{i} (7 \sin u) - \hat{j} (-7 \cos u) + \hat{k} 0 \\ = \langle -7 \sin u, 7 \cos u, 0 \rangle$$

$$\begin{aligned}|\vec{t}_u \times \vec{t}_v| &= \sqrt{7^2 \sin^2 u + 7^2 \cos^2 u} \\ &= \sqrt{7^2(1)} = 7\end{aligned}$$



$$\text{surface area is } \iint_R 7 dA = \int_0^\pi \int_0^8 7 du dv = \boxed{56\pi}$$


We ended here on Friday,
before group quiz

MML #6 (Additional Ex)

call this surface S

Find the area of the surface $z = 5x^2 + 5y^2$ for $0 \leq z \leq 80$.

Sol: S is a paraboloid



Area of S is the surface integral $\iint_S 1 dS$

Projection of S on the xy-plane is

$$R = \{(x, y) : \underbrace{5x^2 + 5y^2 \leq 80}_{x^2 + y^2 \leq 16}\} = \{(r, \theta) : 0 \leq r \leq 4, 0 \leq \theta \leq 2\pi\}$$

Since S is explicitly defined by $z = \underbrace{5x^2 + 5y^2}_{x^2 + y^2}, 0 \leq z \leq 80$,

let $g(x, z) = \underbrace{5x^2 + 5y^2}_z$

$$\text{Then } z_x = 10x, \quad z_y = 10y, \quad \text{so } \sqrt{z_x^2 + z_y^2 + 1} = \sqrt{100(x^2 + y^2) + 1}$$

$$\iint_S 1 dS = \iint_R 1 \sqrt{\underbrace{100(x^2 + y^2) + 1}_{r^2}} dA = \int_0^{2\pi} \int_0^4 \sqrt{100r^2 + 1} r dr d\theta$$

$$\text{inner: } \int_0^4 \sqrt{100r^2 + 1} r dr = \frac{1}{200} \int_{u=1}^{u=1601} u^{\frac{1}{2}} du = \frac{1}{200} \cdot \frac{2}{3} u^{\frac{3}{2}} \Big|_{u=1}^{u=1601} = \frac{1}{300} \left[(1601)^{\frac{3}{2}} - 1 \right]$$

$$du = 200r dr$$

$$\frac{1}{200} du = r dr$$

$$\text{outer: } \int_0^{2\pi} \frac{1}{300} \left[(1601)^{\frac{3}{2}} - 1 \right] d\theta = \boxed{\frac{1}{300} \left[(1601)^{\frac{3}{2}} - 1 \right] 2\pi}$$

8. (10 Pts) If the density of ants on the surface of the paraboloid $z = x^2 + y^2$ ($0 \leq z \leq 4$ m) is given by $f(x, y, z) = 32(x^2 + y^2)$ ($\frac{\text{ants}}{m^2}$), determine the total number of ants on the surface. Note: Focus on expressing the answer as a simplified integral in polar coordinates. Complete the integral if you can, but correct final integration will only be scored 1 point.

Group Quiz Fri

Ref: See Sec 17.6 Example 6, MML # 6

Mass on a surface S is the surface integral $\iint_S (\text{density}) dS = \iint_S f(x, y, z) dS$

Since the surface is explicitly defined by $z = \underbrace{x^2 + y^2}_{\text{let } g(x, z) = x^2 + y^2}, 0 \leq z \leq 4$,

we will use the double integral


THEOREM 17.14 Evaluation of Surface Integrals of Scalar-Valued Functions on Explicitly Defined Surfaces

Let f be a continuous function on a smooth surface S given by $z = g(x, y)$, for (x, y) in a region R . The surface integral of f over S is

$$\iint_S f(x, y, z) dS = \iint_R f(x, y, g(x, y)) \sqrt{z_x^2 + z_y^2 + 1} dA.$$

If $f(x, y, z) = 1$, the surface integral equals the area of the surface.

Sketch of S :



z is from 0 to 4

Set $z = 4$:

$$x^2 + y^2 = 4$$

The projection of S on the xy -plane is $R = \{(x, y) : x^2 + y^2 \leq 4\} = \{(r, \theta) : 0 \leq r \leq 2, 0 \leq \theta \leq 2\pi\}$

$$z_x = 2x, \quad z_y = 2y, \quad \text{so} \quad \sqrt{z_x^2 + z_y^2 + 1} = \sqrt{4x^2 + 4y^2 + 1}$$

$$\iint_S f(x, y, z) dS = \iint_R 32 \underbrace{(x^2 + y^2)}_{r^2} \underbrace{\sqrt{4x^2 + 4y^2 + 1}}_{4r^2} dA$$

write the iterated integral in polar because R is a disk

$$= 32 \int_0^{2\pi} \int_0^2 r^2 \sqrt{4r^2 + 1} r dr d\theta$$

extra

(see Sec 16.3
"Double integrals
in polar coordinates")