


Superunitary regions, generalized associahedra, and





friezes of Dynkin type cluster algebras

Emily Gunawan (UMass Lowell)

Jt. w/ Greg Muller

33rd Meeting on Representation Theory
of Algebras and Related Topics

September 23, 2023




Δ Dynkin diagram \longleftrightarrow [Fomin-Zelevinsky 2001]
 ABCDEFG cluster algebra \mathcal{A}
 of finite type Δ

E.g. Type C_2 Δ : $a \xrightarrow{(2,1)} b$

Choose an orientation of Δ to get a "valued quiver" :

$Q = a \xrightarrow{(2,1)} b$
 \uparrow $\{x, y\}$ \leftarrow initial cluster
 initial (valued) quiver

The exchange graph for type C_2 cluster algebra




The six cluster variables:

- x
- y
- $\frac{1+x^2}{y}$
- $\frac{y+1+x^2}{xy}$
- $\frac{x^2+1+2y+y^2}{x^2y}$
- $\frac{1+y}{x}$

The superunitary region of the C_2 cluster algebra, embedded in \mathbb{R}^2

The six cluster variables,
each set to ≥ 1 :

- $x \geq 1$
- $y \geq 1$
- $\frac{1+x^2}{y} \geq 1$
- $\frac{y+1+x^2}{xy} \geq 1$
- $\frac{x^2+1+2y+y^2}{x^2y} \geq 1$
- $\frac{1+y}{x} \geq 1$




The superunitary region of the A_3 cluster algebra, embedded in \mathbb{R}^3

Initial quiver $\begin{array}{ccc} & 3 & \\ \swarrow & & \searrow \\ 1 & \longrightarrow & 2 \end{array}$

Initial cluster $\{x_1, x_2, x_3\}$

The nine cluster variables,
each set to ≥ 1 :

$$\begin{array}{ccc} x_1 \geq 1 & x_2 \geq 1 & x_3 \geq 1 \\ \frac{x_2+x_3}{x_1} \geq 1 & \frac{x_1+x_3}{x_2} \geq 1 & \frac{x_1+x_2}{x_3} \geq 1 \\ \frac{x_1+x_2+x_3}{x_1x_2} \geq 1 & \frac{x_1+x_2+x_3}{x_2x_3} \geq 1 & \frac{x_1+x_2+x_3}{x_1x_3} \geq 1 \end{array}$$



[Chapoton — Fomin — Zelevinsky 2002]

Let \mathcal{A} be a Dynkin type cluster algebra.

The exchange graph of \mathcal{A} is the 1-skeleton of a polytope called the generalized associahedron of \mathcal{A} .

Faces of the gen. associahedron are indexed by subclusters:

a subcluster is a subset of a cluster

0-face (vertex) \longleftrightarrow cluster $\mathbb{X} = \{x_1, x_2, \dots, x_r\}$
 $\hookrightarrow r := \#$ of vertices of \mathcal{Q}

face \longleftrightarrow subcluster

$(r-1)$ -face (facet) \longleftrightarrow a cluster variable x

1-face (edge) \longleftrightarrow a subcluster \mathbb{X} of size $r-1$ (aka a mutation)

Interior \longleftrightarrow the empty subcluster


Idea Construct a regular CW complex with the same face structure as the generalized associahedron

E.g. for type C_2 cluster algebra:

vertices \leftrightarrow clusters

facets \leftrightarrow cluster variables

interior \leftrightarrow the empty subcluster



Totally positive region

Def \mathcal{A} : a cluster algebra.

- Define a topological space $\mathcal{A}(\mathbb{R}) := \{\text{ring homomorphisms } p: \mathcal{A} \rightarrow \mathbb{R}\}$ with the coarsest topology for which, for all $a \in \mathcal{A}$,
$$f_a: \mathcal{A}(\mathbb{R}) \rightarrow \mathbb{R}$$
the map $p \mapsto p(a)$ is continuous.

- The totally positive region of \mathcal{A} is $\mathcal{A}(\mathbb{R}_{>0}) :=$ the set of ring homomorphisms $p: \mathcal{A} \rightarrow \mathbb{R}$ which send each cluster variable to a positive number.

Fact Given a cluster $\mathcal{X} = \{x_1, x_2, \dots, x_r\}$ in \mathcal{A} ,


we can identify $\mathcal{A}(\mathbb{R}_{>0})$ with the positive orthant $\mathbb{R}_{>0}^r$

via homeomorphism $f_{\mathcal{X}}: \mathcal{A}(\mathbb{R}_{>0}) \cong \mathbb{R}_{>0}^r$
$$p \mapsto (p(x_1), p(x_2), \dots, p(x_r))$$

E.g. $\mathcal{X} = \{x_1, x_2\}$

if $p(x_1) = 1, p(x_2) = 1$ then $f_{\mathcal{X}}(p) = (1, 1)$

if $p(x_1) = 3, p(x_2) = 2$ then $f_{\mathcal{X}}(p) = (3, 2)$



Superunitary region

"bigger than 1"

Main Def The superunitary region of a Dynkin type cluster algebra \mathcal{A} is

$$\mathcal{A}(\mathbb{R}_{\geq 1}) := \left\{ \begin{array}{l} \text{ring homomorphisms } p: \mathcal{A} \rightarrow \mathbb{R} \text{ such that} \\ p(x) \geq 1 \text{ for all cluster variables } x \end{array} \right\} \subset \mathcal{A}(\mathbb{R}_{> 0})$$

Totally positive region

Given a cluster \mathbb{X} , use homeomorphism $f_{\mathbb{X}}: \mathcal{A}(\mathbb{R}_{> 0}) \xrightarrow{\cong} \mathbb{R}_{> 0}^r$

to embed $\mathcal{A}(\mathbb{R}_{\geq 1})$ into $\mathbb{R}_{> 0}^r$:
superunitary region

Totally positive region


Positive orthant

Set each cluster variable ≥ 1

(positive Laurent polynomial)

E.g. Embedding of type C_2 $\mathcal{A}(\mathbb{R}_{\geq 1})$ into \mathbb{R}^2

using $a \xrightarrow{(2,1)} b$:
 $\mathbb{X} = \{x, y\}$



$$\mathbb{X} := \{x, y\}$$

In this embedding $f_{\mathbb{X}}$,

\mathbb{P} where $\mathbb{P}(x) = 3, \mathbb{P}(y) = 2$

$$\mathbb{P}\left(\frac{1+x^2}{y}\right) = \frac{1+9}{2} = 5$$

$$\mathbb{P}\left(\frac{y+1+x^2}{xy}\right) = \frac{2+1+9}{6} = 2$$


$$\mathbb{P}\left(\frac{x^2+1+2y+y^2}{x^2y}\right) = \frac{9+1+4+4}{18} = 1$$

$$\mathbb{P}\left(\frac{1+y}{x}\right) = \frac{1+2}{3} = 1$$

$$f_{\mathbb{X}} \mapsto (3, 2) \in \mathbb{R}^2$$

(an extreme point)

all integer



$$\mathbb{X} := \{x, y\}$$

In this embedding $f_{\mathbb{X}}$,


p where $p(x)=2, p(y)=3 \xrightarrow{f_{\mathbb{X}}} (2,3) \in \mathbb{R}^2$
(an interior point)

$$p\left(\frac{1+x^2}{y}\right) = \frac{1+4}{3} = \frac{5}{3}$$

$$p\left(\frac{y+1+x^2}{xy}\right) = \frac{3+1+4}{6} = \frac{8}{6} = \frac{4}{3}$$

$$p\left(\frac{x^2+1+2y+y^2}{x^2y}\right) = \frac{4+1+6+9}{12} = \frac{20}{12} = \frac{5}{3}$$

$$p\left(\frac{1+y}{x}\right) = \frac{1+3}{2} = 2$$



$$\mathbb{X} := \{x, y\}$$

In this embedding $f_{\mathbb{X}}$,

\mathbb{P} where $\mathbb{P}(x) = 1, \mathbb{P}(y) = 1$

$$\mathbb{P}\left(\frac{1+x^2}{y}\right) = 2$$


$$\mathbb{P}\left(\frac{y+1+x^2}{xy}\right) = 3$$

$$\mathbb{P}\left(\frac{x^2+1+2y+y^2}{x^2y}\right) = 5$$

$$\mathbb{P}\left(\frac{1+y}{x}\right) = 2$$

$f_{\mathbb{X}} \mapsto (1,1) \in \mathbb{R}^2$
(an extreme point)

all integer



Thm A If \mathcal{A} is a Dynkin type cluster algebra, the superunitary region $\mathcal{A}(\mathbb{R}_{\geq 1})$ is a regular CW complex which is cellular homeomorphic to the generalized associahedron.

Subcluster face indexed by subcluster \mathbb{X} is $\{p: \mathcal{A} \rightarrow \mathbb{R} \text{ s.t. } p(a) = 1 \text{ iff } a \in \mathbb{X}\}$

- k-face \longleftrightarrow a subcluster \mathbb{X} of size $r-k$
- (r-1)-face (facet) \longleftrightarrow a cluster variable x
- 1-face (edge) \longleftrightarrow a subcluster \mathbb{X} of size $r-1$ (aka a mutation)
- 0-face (vertex) \longleftrightarrow a cluster $\mathbb{X} = \{x_1, x_2, \dots, x_r\}$

- Boundary of $\mathcal{A}(\mathbb{R}_{\geq 1}) = \bigsqcup_{\mathbb{X} \text{ nonempty}} \text{subcluster face indexed by } \mathbb{X}$
- Interior of $\mathcal{A}(\mathbb{R}_{\geq 1}) = \{p: \mathcal{A} \rightarrow \mathbb{R} \text{ where } p(x) > 1 \forall \text{ cl. var } x\}$
indexed by the empty subcluster

Cor $\mathcal{A}(\mathbb{R}_{\geq 1})$ is closed and bounded.

Pf The generalized associahedron is a polytope. \blacksquare

An application of superunitary regions:

A uniform proof of a previously open conjecture that there are finitely many positive integral friezes, for each Dynkin type.

Def The set of frieze points is

$$\mathcal{A}(\mathbb{Z}_{\geq 1}) := \left\{ \text{ring homomorphisms } p: \mathcal{A} \rightarrow \mathbb{R} \text{ such that } \begin{array}{l} p(x) \in \mathbb{Z}_{\geq 1} \text{ for all cluster variables } x \end{array} \right\} \subset \mathcal{A}(\mathbb{R}_{\geq 1})$$

superunitary region

Cor If \mathcal{A} is a Dynkin type cluster algebra, the set of frieze points is finite.

Pf The homeomorphism $\mathcal{A}(\mathbb{R}_{>0}) \simeq \mathbb{R}_{>0}^r$ totally positive region

restricts to

$$\mathcal{A}(\mathbb{Z}_{\geq 1}) \hookrightarrow \mathbb{Z}_{\geq 1}^r$$

positive orthant

pos integral points


and


$$\mathcal{A}(\mathbb{Z}_{\geq 1}) \hookrightarrow \mathcal{A}(\mathbb{R}_{\geq 1})$$

superunitary region


Since $\mathcal{A}(\mathbb{R}_{\geq 1})$ is bounded, the set of frieze points is finite. \square

What do we mean by positive integral friezes (in this talk)?


Def Q a (valued) Dynkin quiver, e.g. $Q =$

 (type A_4)
 Build the "repetition quiver" $\mathbb{Z}Q$




A positive integral frieze is a function $\mathbb{Z}Q \rightarrow \mathbb{Z}_{\geq 1}$ satisfying ...

• [Conway-Coxeter 1970s, type A] for each , we have $ad - bc = 1$

E.g. a type $A_3 \mathbb{Z}_{\geq 1}$ -frieze



• [Caldero-Chapoton 2006 and Assem-Reutenaur-Smith 2010, in general]

for each , $ad - b_1 \dots b_k = 1$

→ friezes \leftrightarrow cluster algebras

Fact If Q is Dynkin, $\mathcal{A}(\mathbb{Z}_{\geq 1})$ $\xleftrightarrow{|-|}$ $\mathbb{Z}_{\geq 1}$ -friezes of Q
frieze points

Thm B If Q is Dynkin, there are finitely many $\mathbb{Z}_{\geq 1}$ -friezes of Q .

Pf Earlier we said $\mathcal{A}(\mathbb{Z}_{\geq 1})$ is a finite set.

History of proofs, by type	Techniques
✓ Type A Conway-Coxeter 1970s	Polygon triangulations
✓ BCD, G_2 Fontaine-Plamondon 2014	Type D triangulations (once-punctured polygon)
✓ E_6, F_4 Cuntz-Plamondon 2018	$\{E_6 \text{ friezes}\} \hookrightarrow \{2\text{-friezes of height } 3\}$
✓ E_7, E_8 G. - Muller 2022 Conjecture for E_7, E_8 was open until	Uniform proof for all types using compactness of the superunitary region

Conjecture C Any embedding of the superunitary region $\mathcal{A}(\mathbb{R}_{\geq 1})$ is contained in the convex hull of the extreme points.

Dynkin Type	# of Positive Integral Friezes	# of Unitary Friezes = (# of Clusters) = (# of extreme points)
A_n	$\frac{1}{n+2} \binom{2n+2}{n+1}$	$\frac{1}{n+2} \binom{2n+2}{n+1}$
B_n	$\sum_{m=1}^{\sqrt{n+1}} \binom{2n-m^2+1}{n}$	$\binom{2n}{n}$
C_n	$\binom{2n}{n}$	$\binom{2n}{n}$
D_n	$\sum_{m=1}^n d(m) \binom{2n-m-1}{n-m}$	$\frac{3n-2}{n} \binom{2n-2}{n-1}$
E_6	868	833
E_7	Open (conjectured: 4400)	4160
E_8	Open (conjectured: 26952)	25080
F_4	112	105
G_2	9	8

To prove these ...
 • prove Conjecture, or
 • prove # of $\mathbb{Z}_{\geq 2}$ -valued friezes of type E_7 & E_8 are 0 and 4

$d(m) :=$ the number of divisors of m

TABLE 1. Counts of positive integral friezes

Thm D There is a frieze point
in the interior of $\mathcal{A}(\mathbb{R}_{\geq 1})$
iff

Q is a union of ...

- type D_n , n not prime

- type E_8

- type B_n , $\sqrt{n+1} \in \mathbb{Z}_{\geq 2}$

- type G_2

(assuming conjectured # of
type E_7, E_8 positive integral friezes)

E.g. In rank 2, there is
a frieze point in
the interior of $\mathcal{A}(\mathbb{R}_{\geq 1})$
iff Q is of type G_2



Quiver	Dynkin type	Superunitary region	Inequalities
$\circ \quad \circ$	$A_1 \times A_1$		$\begin{aligned} x_1 &\geq 1 \\ 2 &\geq x_2 \\ 2 &\geq x_1 \\ x_2 &\geq 1 \end{aligned}$
$\circ \longrightarrow \circ$	A_2		$\begin{aligned} x_1 &\geq 1 \\ x_1 + 1 &\geq x_2 \\ x_1 + x_2 + 1 &\geq x_1 x_2 \\ x_2 + 1 &\geq x_1 \\ x_2 &\geq 1 \end{aligned}$
$\circ \xrightarrow{(2,1)} \circ$	B_2 or C_2		$\begin{aligned} x_1 &\geq 1 \\ x_1^2 + 1 &\geq x_2 \\ x_1^2 + x_2 + 1 &\geq x_1 x_2 \\ x_2^2 + 2x_2 + x_1^2 + 1 &\geq x_1^2 x_2 \\ x_2 + 1 &\geq x_1 \\ x_2 &\geq 1 \end{aligned}$
$\circ \xrightarrow{(3,1)} \circ$	G_2		$\begin{aligned} x_1 &\geq 1 \\ \frac{x_1^3 + 1}{x_2} &\geq 1 \\ \frac{x_1^3 + x_2 + 1}{x_1} &\geq 1 \\ \frac{x_1^6 + 3x_2x_1^3 + 2x_1^3 + x_2^3 + 3x_2^2 + 3x_2 + 1}{x_2^2 x_1^3} &\geq 1 \\ \frac{x_2^3 + 3x_2^2 + 3x_2 + x_1^3 + 1}{x_2 x_1^3} &\geq 1 \\ \frac{x_1^3 + x_2^2 + 2x_2 + 1}{x_2 x_1^2} &\geq 1 \\ \frac{x_2 + 1}{x_1} &\geq 1 \\ x_2 &\geq 1 \end{aligned}$

THANK
YOU




FIGURE 1. The superunitary regions of types A_2 , B_2/C_2 , and G_2 (embedded in $\mathbb{R}_{>0}^2$)